

The short-term and long-term effects of cannabis on cognition: recent advances in the field

Emese Kroon^{1,2}, Lauren Kuhns^{1,2} and Janna Cousijn^{1,2}

The aim of this review is to discuss the most recent evidence for the short-term and long-term effects of cannabis on cognition. The evidence that cannabis intoxication is associated with short-term impairment across several basal cognitive domains, including learning and (episodic) memory, attentional control, and motor inhibition is increasing. However, evidence regarding the effects of long-term heavy cannabis use on cognition remains equivocal. Cannabis research suffers from difficulties in measuring cannabis exposure history, poor control over potential subacute effects, and heterogeneity in cognitive measures and sample composition. Multidisciplinary collaborations and investment in studies that help overcome these difficulties should be prioritized.

Addresses

¹ Neuroscience of Addiction (NofA) Lab, Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands

² The Amsterdam Brain and Cognition Center (ABC), University of Amsterdam, Amsterdam, The Netherlands

Corresponding author: Kroon, Emese (e.kroon@uva.nl)

Current Opinion in Psychology 2021, 38:49–55

This review comes from a themed issue on **Cannabis**

Edited by **Ken C Winters** and **Joel Mader**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 16th July 2020

<https://doi.org/10.1016/j.copsyc.2020.07.005>

2352-250X/© 2020 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Recent global changes in cannabis legislation parallel increases in use and decreases in harm perception [1,2]. Yet, there is still little conclusive evidence on the effects of cannabis use. This review specifically focuses on the effects of cannabis use on cognition. Cognition encompasses our thoughts and shapes our behaviour, and refers to distinct but partially overlapping processes such as learning, memory, attention, inhibition, decision-making, and emotion regulation. Cannabis contains over a hundred different cannabinoids including Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD [3]). Although the mechanisms are unclear, cannabinoids like THC and CBD potentially affect cognition through interactions with the endogenous cannabinoid system in the brain [4]. This system in-turn regulates many other

neurotransmitter systems including the dopamine system often implicated in substance use disorders (SUD [5]). Moreover, like in other SUDs, the development of Cannabis Use Disorder (CUD) may also be related to pre-existing cognitive deficits [6]. Given the rapidly developing evidence base, we will discuss the most recent evidence for the effects of cannabis intoxication (short-term) and heavy cannabis use (almost daily use, long-term) on cognition (Table 1). We thereby start with basal cognitive functions, moving towards more complex cognitive functions and the role of affective processes therein.

Cannabis and cognition: current knowledge and recent advances

Learning and memory

Cannabis intoxication impairs learning and memory in a dose-dependent manner, although significant individual differences exist [7,8,9^{*}]. Studies in heavy cannabis users are less consistent, but learning and immediate recall deficits are most commonly reported in active cannabis users [10^{**}]. A recent longitudinal study [11^{*}] in adolescent cannabis users suggests a causal link between cannabis exposure and immediate, but not delayed recall in an episodic memory task. Furthermore, another recent study showed that trial-by-trial verbal learning rates were slower in cannabis users compared to controls, and these learning rates were associated with altered functionality of the parahippocampal gyrus, thalamus and midbrain regions [12^{*}]. While altered feedback processing may play a role in learning deficits observed in alcohol and other substance users, this may not necessarily be the case in cannabis users [13]. Furthermore, impairments may not be relegated to only memory of real experiences. Kloft *et al.* [14^{*}] showed that cannabis intoxication increased susceptibility to false memory, an effect that appeared most prominent at immediate compared to delayed recall.

Subacute intoxication effects likely contribute to the described effects in cannabis users. The effects of cannabis on memory performance and related alterations in brain activity fade with abstinence [10^{**}]. In line with this, working memory performance and functionality of the underlying brain network was only found to be impaired in individuals with a positive urine screen for THC [15^{*}]. Despite the heterogeneous and potential timebound nature of the observed deficits, cannabis use-related learning and memory problems could seriously impact daily functioning of heavy cannabis users, including performance in school or at work. A combination of psychological, neurological, and neurobiological

Table 1

Summary of current evidence for short-term and long-term effects of cannabis on cognition

Domain	Short-term effects		Long-term effects		Suggested reading	
	Evidence	Potential moderators	Evidence	Potential moderators	Reviews	Recent evidence
Learning and memory	Sufficient evidence that THC/cannabis impairs (non)-verbal learning and episodic memory. Limited evidence for impairment of other types of learning and memory.	Dose ↑ Early onset ↑ Heavy history ↓ Low THC:CBD ratio ↓	Sufficient evidence for impairments in current heavy users. Insufficient evidence for lasting effects after abstinence. Indications of (partial) recovery.	Subacute THC/cannabis effects ↑ Early onset ↑ Heavy history ↑ Comorbid mental health issues ↑	[7,8,10**,16**]	[11*,12*,13,14*,36,37*]
Working memory	Inconsistent evidence that THC/cannabis impairs working memory.	–	Inconsistent evidence for long-term working memory deficits in current heavy users. Limited evidence for recovery after abstinence.	Subacute THC/cannabis effects ↑ Heavy history ↑ Early onset ↑ Task complexity ↑	[7,8,10**]	[15*]
Attentional control	Sufficient evidence that THC/cannabis impairs attentional control.	Dose ↑ Heavy history ↓	Sufficient evidence for impairments in sustained and divided attention in current heavy users. Insufficient evidence for lasting effects after abstinence. Indications of (partial) recovery.	Subacute THC/cannabis effects ↑ Early onset ↑ Heavy history ↑	[29,42**,52]	[9*]
Motor inhibition	Sufficient evidence that THC/cannabis impairs inhibition of ongoing responses (stop-signal task). Inconsistent results with other inhibition tasks.	Dose ↑	Limited and inconsistent evidence for impairments in current heavy users.	–	[29,42**,52]	[9*]
Cognitive biases	Limited evidence for cannabis-related approach bias and attentional bias.	–	Sufficient evidence for attentional bias, but insufficient evidence for approach bias in current heavy users. No evidence to support or refute lasting effects after abstinence.	Heavy history ↑ CUD severity ↑ THC ↑ Craving ↑	[24**,53]	[22,23,26*]
Emotion processing	Consistent , but limited evidence that THC/cannabis impairs emotion recognition, particularly for negative emotions.	Low THC:CBD ratio ↓	Limited evidence for impaired emotion identification/recognition in current heavy users. No evidence to support or refute lasting effects after abstinence.	–	–	[41*]
Decision making	Insufficient evidence that THC/cannabis impairs decision-making.	–	Insufficient and inconsistent evidence for impairments in current heavy users.	Cognitive subdomain	[29,43,53]	[11*]

This table is an adaptation and update of the table presented in Kroon *et al.* [31], focussing on the existing knowledge and most recent evidence for short-term and long-term effects of cannabis on cognition. The short-term effects column includes results from intoxication studies, while the long-term effects column includes evidence for the effects of longer periods of heavy (near daily) cannabis use on cognition.

THC = Δ 9-tetrahydrocannabinol; CBD = cannabidiol.

research [16**] is crucial to further elucidate the apparent complexity of mechanisms underlying the effects of cannabis on memory.

Attention

Similar to learning and memory, cannabis intoxication consistently results in a THC-dose-dependent reduction of the capacity to orient attention towards task-relevant stimuli [17–19]. In heavy compared to occasional cannabis users, tolerance to the acute effect of cannabis on attentional control was related to reduced responsiveness of the reward system after intoxication [20*]. This may relate to the general tolerance to cognitive impairments by cannabis intoxication often observed in heavy users [7,8,17,18,21]. Heavy cannabis users also develop an attentional bias towards cannabis and related objects that may interfere with other attentional processes [e.g. Ref. [22], but see Ref. [23]]. Although effect sizes were small, a recent meta-analysis showed evidence for an attentional bias towards cannabis-related words and pictures in heavy cannabis users [24**]. Attentional bias has been linked to the severity of CUD [25] and might reflect an involuntary early perceptual bias, supported by increased amplitude and earlier peak of the N1 component in response to distracting cannabis stimuli [26*].

Inhibition

Cannabis use, and drug use in general, has often been associated with poor inhibitory control. With regards to motor inhibition, cannabis intoxication consistently and dose dependently reduces the ability to inhibit an ongoing motor response, as measured with the stop-signal task (e.g. Refs. [27,28]). In contrast, inhibition before a response is initiated, as measured with the go/no-go task, may not be impaired by intoxication [28]. Findings on the effects of heavy cannabis use on motor inhibition are less consistent [29]. However, aside from potential problems caused by impairments in motor control due to cannabis intoxication [30], motor inhibition might not well-reflect the daily life inhibition problems present in most substance users. Indeed, slower proactive inhibitory control-related processes, such as those measured with the classical Stroop were found to relate to cannabis craving [23].

Decision-making

More complex cognitive functions such as decision-making heavily rely on the integrity of the basal cognitive functions discussed above and deficits in any of those might in turn result in risky decisions like substance use. The complexity of the processes involved may explain the inconsistent findings on the effects of cannabis intoxication and heavy use on decision-making [29,31]. Nonetheless, progress has been made and recent studies provide new insight into how heavy cannabis use and the context in which decisions are made affect risky decision-making. For example, a recent study on financial delay discounting (preferring immediate small rewards over

delayed bigger rewards) observed a positive relationship between increased delay discounting and frequency of cannabis use [32]. Interestingly, Gilman *et al.* [33] found that heavy cannabis using adolescents compared to controls differed on risk taking in the social, safety, and ethical domains, but not the financial domain. In general, risky decision-making in heavy cannabis users seems associated with increased sensitivity to immediate gain accompanied by decreased loss sensitivity [34,35].

The importance of context and emotion

The previously discussed findings highlight the need for a more fine-grained investigation of cognitive subprocesses and their interactions, as well as the importance of the context in which cognition is measured. While cannabis use by a popular peer may bias decision-making in an occasional user, for individuals with a CUD, decision-making may be particularly compromised when confronted with cannabis-related cues. As with attentional bias, cannabis-related cues may also activate an approach bias towards cannabis in heavy cannabis users [25]. Moreover, acute stress may influence cognitive performance. For example, acute stress affects prospective memory performance in both heavy cannabis users and controls, but the effects are larger in heavy cannabis users [36]. On the other hand, increased working memory capacity seems to protect heavy cannabis users from craving under stressful circumstances [37*]. Taken together, potential cognitive deficits in heavy cannabis users may manifest themselves depending on contextual factors.

The impact of cannabis use on emotion processing is an important factor to consider herein. Although data is limited, cannabis intoxication may negatively affect emotion recognition [38]. This seems to be most apparent for negative emotions and appears to be related to reduced brain activity in reward and cognitive control related brain areas when presented with negative faces [39,40]. A recent study focusing on gender differences identified complex interactions between gender and cannabis use patterns in relation to the early processing of emotional stimuli (EEG, ERP: P1 and P3 [41*]). This highlights the general importance of assessing gender differences in the effects of cannabis use. This is a particularly relevant issue in the domain of emotion processing research because of the high rates of comorbidity between cannabis use and disorders associated with emotion processing (e.g. anxiety) and the commonly reported gender difference in the prevalence of these disorders.

Field wide difficulties and future directions

Aside from the classic confounders such as polysubstance use and comorbid mental health problems, as well as a lack of longitudinal data limiting our understanding of the causal relationship between cannabis and cognition, cannabis research is facing significant difficulties which have been brought to attention by the majority of recent

reviews on the topic [10^{**},24^{**},42^{**},43]. While overcoming these difficulties is of utmost importance, clear solutions are still lacking.

First, the vast majority of studies on the long-term effects of heavy cannabis use on cognition share one confounding factor: the abstinence period. Studies show that THC metabolites are detectable in the plasma of heavy cannabis users for over a week [44] and even longer detectability is possible due to THC's lipophilic characteristics [45]. In line with this, cannabis-use-dependent neurocognitive impairments can be detected for as long as 28 days after cessation [46]. Hence, studies in current heavy cannabis users struggle to differentiate subacute from long-term effects. Although this confound should be acknowledged and more wide-spread assessment of THC metabolites is warranted, subacute effects should not always be seen as a problem in itself. After all, the mix of acute, subacute, and long-term effects represent what a current heavy cannabis user is dealing with in daily life. Nevertheless, more knowledge of the potential for recovery after abstinence and the role of CUD severity in recovery is needed.

Second, problems with quantifying use are often reported and pose a true problem for comparability across studies. Variable definitions of heavy cannabis use and the lack of standard cannabis units are recurrent problems. While both problems might reflect semantics, and defining categories for frequency and heaviness of use might indeed primarily require discussion, developing a standard unit is extremely complicated. Recently, attempts were made to develop a standard unit of cannabis [47,48^{**}], but the complexity and variability in cannabis products and routes of administration hampers practicality. Cannabis contains over a hundred different types of cannabinoids and the THC:CBD ratio differs significantly between region and even between batches [49]. Poor knowledge about exposure history in most studies complicates research even further. To improve our knowledge base, accessible and more reliable methods to quantify cannabis use are needed. However, even then, research in most countries heavily relies on changes in local legislation to allow for these methods to be used.

Third, there are methodological problems that plague comparability in systematic reviews and meta-analyses. While increasing the amount of research will increase the power of these types of reviews, studies are rarely replicated and the variability between measures to assess the same cognitive construct remains a problem [24^{**},42^{**},43]. An increase in power will not reflect an increase in knowledge when this heterogeneity problem is not solved. In line with this, it remains important to be aware of the risks of assuming that similar tasks measure

the same construct like is often done when aggregating results from stop-signal and go/no-go task [50].

Finally, it may be that the effects of heavy cannabis use on cognition are indeed mixed. The same dose of THC may result in impairments in some, while leading to improvement in others [51]. These individual differences are likely to depend on a large variety of moderating factors including THC:CBD ratio, differences in THC metabolism, poly-substance use, severity of cannabis dependence, age of onset, gender, and mental health. In turn, the combined effects of these factors might vary with the context under which cannabis is consumed and cognition is assessed.

Conclusion

The rapid increase of research into cannabis and its effects on cognition has provided us with answers as well as questions. While there is increasing evidence that cannabis intoxication negatively affects basal cognitive functions like episodic memory, attentional control, and motor inhibition, results on the long-term effects of heavy cannabis use, and potential recovery after abstinence, remain equivocal for most cognitive domains. Despite a slow start, cannabis research is breaking ground. Nevertheless, field-wide difficulties in quantification, methods of measuring cognitive constructs, and the influence of subacute effects seriously hamper the road ahead and require attention now. Multidisciplinary collaboration and investment in studies that solve these problems should be prioritized.

Conflict of interest statement

Nothing declared.

Acknowledgements and funding

This review was supported by grant 1R01 DA042490-01A1 from the National Institute on Drug Abuse/National Institutes of Health.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. UNODC: **Cannabis and hallucinogens**. *World Drug Report 2019*. 2019:1-71.
2. SAMHSA: **Key Substance Use and Mental Health Indicators in the United States: Results From the 2017 National Survey on Drug Use and Health**. (HHS Publication No. PEP19-5068, NSDUH Series H-54). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration; 2018.
3. Chandra S, Radwan MM, Majumdar CG, Church JC, Freeman TP, ElSohly MA: **New trends in cannabis potency in USA and Europe during the last decade (2008–2017)**. *Eur Arch Psychiatry Clin Neurosci* 2019, **269**:5-15 <http://dx.doi.org/10.1007/s00406-019-00983-5>.
4. Russo EB: **Beyond cannabis: plants and the endocannabinoid system**. *Trends Pharmacol Sci* 2016, **37**:594-605 <http://dx.doi.org/10.1016/j.tips.2016.04.005>.

5. Covey DP, Mateo Y, Sulzer D, Cheer JF, Lovinger DM: **Endocannabinoid modulation of dopamine neurotransmission.** *Neuropharmacology* 2017, **124**:52-61 <http://dx.doi.org/10.1016/j.neuropharm.2017.04.033>.
6. Bickel WK, Mellis AM, Snider SE, Athamneh LN, Stein JS, Pope DA: **21st century neurobehavioral theories of decision making in addiction: review and evaluation.** *Pharmacol Biochem Behav* 2018, **164**:4-21 <http://dx.doi.org/10.1016/j.pbb.2017.09.009>.
7. Schoeler T, Bhattacharyya S: **The effect of cannabis use on memory function: an update.** *Subst Abuse Rehabil* 2013, **4**:11-27 <http://dx.doi.org/10.2147/SAR.S25869>.
8. Ranganathan M, D'Souza DC: **The acute effects of cannabinoids on memory in humans: a review.** *Psychopharmacology (Berl)* 2006, **188**:425-444 <http://dx.doi.org/10.1007/s00213-006-0508-y>.
9. Petker T, Owens MM, Amlung MT, Oshri A, Sweet LH, Mackillop J: **Cannabis involvement and neuropsychological performance: Findings from the human connectome project.** *J. Psychiatry Neurosci* 2019, **44**:414-422 <http://dx.doi.org/10.1503/jpn.180115>.
- 9]Petker T, Owens MM, Amlung MT, Oshri A, Sweet LH, Mackillop J: Cannabis involvement and neuropsychological performance: Findings from the human connectome project. *J. Psychiatry Neurosci* 2019, **44**:414-422. <https://doi.org/10.1503/jpn.180115>. A study assessing associations between cannabis use (recent use, lifetime use, cannabis use disorder, and onset of use) and neuropsychological performance (eight different domains) in a large group of young adults ($N=11121$; part of human connectome project). Results show that positive THC screening was associated with worse episodic memory and reduced processing speed. On the other hand, positive CUD status was associated with lower fluid intelligence.
10. Blest-Hopley G, Giampietro V, Bhattacharyya S: **A systematic review of human neuroimaging evidence of memory-related functional alterations associated with cannabis use complemented with preclinical and human evidence of memory performance alterations.** *Brain Sci* 2020, **10**:102 <http://dx.doi.org/10.3390/brainsci10020102>.
- 10]Blest-Hopley G, Giampietro V, Bhattacharyya S: A systematic review of human neuroimaging evidence of memory-related functional alterations associated with cannabis use complemented with preclinical and human evidence of memory performance alterations. *Brain Sci* 2020, **10**:102. <https://doi.org/10.3390/brainsci10020102>. An extensive systematic review of the evidence for a relationship between memory and cannabis use, including human neuroimaging studies, preclinical behavioural studies, and human behavioural studies. An informative distinction between the evidence for the effects on adolescents and adults is made and problems in the literature are extensively discussed.
11. Duperrouzel JC, Hawes SW, Lopez-Quintero C, Pacheco-Colón I, Coxé S, Hayes T, Gonzalez R: **Adolescent cannabis use and its associations with decision-making and episodic memory: Preliminary results from a longitudinal study.** *Neuropsychology* 2019, **33**:701-710 <http://dx.doi.org/10.1037/neu0000538>.
- 11]Duperrouzel JC, Hawes SW, Lopez-Quintero C, Pacheco-Colón I, Coxé S, Hayes T, Gonzalez R: Adolescent cannabis use and its associations with decision-making and episodic memory: Preliminary results from a longitudinal study. *Neuropsychology* 2019, **33**:701-710. <https://doi.org/10.1037/neu0000538>. A longitudinal study presenting preliminary results on the association between adolescent cannabis use ($N = N401$) and decision-making, as well as episodic memory. Results show that there were cross-sectional associations between cannabis use and poorer decision-making and episodic memory. These decision-making deficits were not predictive of use at 1-year-follow-up, suggesting decision-making deficits as measured with this paradigm might not relate to escalation of use. However, an increase in cannabis use at follow-up was found to be associated with decreased immediate recall over that period.
12. Blest-Hopley G, O'Neill A, Wilson R, Giampietro V, Bhattacharyya S: **Disrupted parahippocampal and midbrain function underlie slower verbal learning in adolescent-onset regular cannabis use.** *Psychopharmacology (Berl)* 2019:1-17 <http://dx.doi.org/10.1007/s00213-019-05407-9>.
- 12]Blest-Hopley G, O'Neill A, Wilson R, Giampietro V, Bhattacharyya S: Disrupted parahippocampal and midbrain function underlie slower verbal learning in adolescent-onset regular cannabis use. *Psychopharmacology (Berl)* 2019, **1-17**. <https://doi.org/10.1007/s00213-019-05407-9>. A fMRI study focussed on the brain correlates of verbal learning in cannabis users ($N = N42$). Trial-by-trial analysis of the verbal learning task showed that cannabis users had slower verbal learning rates than non-using controls, which in turn was found to be associated with disrupted brain activity in the midbrain, parahippocampal gyrus and thalamus. These results point towards the importance of studying learning processes, not only learning outcomes.
13. Aloï J, Blair KS, Crum KI, Bashford-Largo J, Zhang R, Lukoff J, Carollo E, White SF, Hwang S, Filbey FM *et al.*: **Alcohol use disorder, but not cannabis use disorder, symptomatology in adolescents is associated with reduced differential responsiveness to reward versus punishment feedback during instrumental learning.** *Biol Psychiatry Cogn Neurosci Neuroimaging* 2020, **5**:610-618 <http://dx.doi.org/10.1016/j.bpsc.2020.02.003>.
14. Kloft L, Otgaar H, Blokland A, Monds LA, Toennes SW, Loftus EF, Ramaekers JG: **Cannabis increases susceptibility to false memory.** *Proc Natl Acad Sci U S A* 2020, **117**:4585-4589 <http://dx.doi.org/10.1073/pnas.1920162117>.
- 14]Kloft L, Otgaar H, Blokland A, Monds LA, Toennes SW, Loftus EF, Ramaekers JG: Cannabis increases susceptibility to false memory. *Proc Natl Acad Sci U S A* 2020, **117**:4585-4589. <https://doi.org/10.1073/pnas.1920162117>. A double-blind, randomized, placebo-controlled study ($N=N64$) assessing the effects of THC on false memory. Result show that cannabis intoxication increased false memory formation and that these effects were most prominent in the acute-intoxication phase.
15. Owens MM, McNally S, Petker T, Amlung MT, Balodis IM, Sweet LH, MacKillop J: **rinary tetrahydrocannabinol is associated with poorer working memory performance and alterations in associated brain activity.** *s associated with poorer working memory performance and alterations in associated brain activity* *Neuropsychopharmacology* 2019, **44**:613-619 <http://dx.doi.org/10.1038/s41386-018-0240-4>.
- 15]Owens MM, McNally S, Petker T, Amlung MT, Balodis IM, Sweet LH, MacKillop J: **rinary tetrahydrocannabinol is associated with poorer working memory performance and alterations in associated brain activity.** *s associated with poorer working memory performance and alterations in associated brain activity* *Neuropsychopharmacology* 2019, **44**:613-619. <https://doi.org/10.1038/s41386-018-0240-4>. A fMRI study assessing the association between cannabis use and working memory (WM) performance and WM related brain activity in a large group of young adults ($N = N1038$; part of human connectome project). Results show that positive THC screening was associated with worse WM performance and that reduced brain activity in WM-related regions in combination with increased activity in WM-unrelated regions mediated this relationship. Moreover, both WM performance and related brain activity were not associated with history of cannabis use, suggesting short-term rather than long-term effects of cannabis on WM.
16. Prini P, Zamberletti E, Manenti C, Gabaglio M, Parolaro D, Rubino T: **Neurobiological mechanisms underlying cannabis-induced memory impairment.** *Eur Neuropsychopharmacol* 2020, **36**:181-190 <http://dx.doi.org/10.1016/j.euroneuro.2020.02.002>.
- 16]Prini P, Zamberletti E, Manenti C, Gabaglio M, Parolaro D, Rubino T: **Neurobiological mechanisms underlying cannabis-induced memory impairment.** *Eur Neuropsychopharmacol* 2020, **36**:181-190. <http://dx.doi.org/10.1016/j.euroneuro.2020.02.002>. review on cannabis-induced memory impairment in which animal models are discussed to explain potential neurobiological mechanisms underlying these impairments.
17. Theunissen EL, Kauert GF, Toennes SW, Moeller MR, Sambeth A, Blanchard MM, Ramaekers JG: **Neurophysiological functioning of occasional and heavy cannabis users during THC intoxication.** *Psychopharmacology (Berl)* 2012, **220**:341-350 <http://dx.doi.org/10.1007/s00213-011-2479-x>.
18. Ramaekers JG, Kauert G, Theunissen EL, Toennes SW, Moeller MR: **Neurocognitive performance during acute THC intoxication in heavy and occasional cannabis users.** *J Psychopharmacol* 2009, **23**:266-277 <http://dx.doi.org/10.1177/0269881108092393>.
19. D'Souza DC, Ranganathan M, Braley G, Gueorguieva R, Zimolo Z, Cooper T, Perry E, Krystal J: **Blunted psychotomimetic and amnesic effects of Δ -9-tetrahydrocannabinol in frequent users of cannabis.** *Neuropsychopharmacology* 2008, **33**:2505-2516 <http://dx.doi.org/10.1038/sj.npp.1301643>.
20. Mason NL, Theunissen EL, Hutten NRPW, Tse DHY, Toennes SW, Jansen JFA, Stiers P, Ramaekers JG: **Reduced responsiveness of the reward system is associated with tolerance to cannabis impairment in chronic users.** *Addict Biol* 2019:e12870 <http://dx.doi.org/10.1111/adb.12870>.
- 20]Mason NL, Theunissen EL, Hutten NRPW, Tse DHY, Toennes SW, Jansen JFA, Stiers P, Ramaekers JG: Reduced responsiveness of the reward system is associated with tolerance to cannabis impairment in chronic users. *Addict Biol* 2019, **e12870**. <https://doi.org/10.1111/adb.12870>. A double-blind, randomized, placebo-controlled, cross-over study ($N = N24$) in which the acute effects of THC on behavioural (sustained attention and subjective high) and brain-related outcomes

(7T fMRI and spectroscopy) were assessed. Results show THC-induced alterations in the reward circuit in occasional cannabis users, associated with increased subjective high and reduced sustained attention. However, these alterations were absent in heavy cannabis users, suggesting tolerance related reduced responsiveness of the reward system in heavy cannabis users.

21. Schwoppe DM, Bosker WM, Ramaekers JG, Gorelick DA, Huestis MA: **Psychomotor performance, subjective and physiological effects and whole blood 9-tetrahydrocannabinol concentrations in heavy, chronic cannabis smokers following acute smoked cannabis.** *J Anal Toxicol* 2012, **36**:405-412 <http://dx.doi.org/10.1093/jat/bks044>.
22. Alcorn JL, Marks KR, Stoops WW, Rush CR, Lile JA: **Attentional bias to cannabis cues in cannabis users but not cocaine users.** *Addict Behav* 2019, **88**:129-136 <http://dx.doi.org/10.1016/j.addbeh.2018.08.023>.
23. Van Kampen AD, Cousijn J, Engel C, Rinck M, Dijkstra BAG: **Addictive behaviors attentional bias, craving and cannabis use in an inpatient sample of adolescents and young adults diagnosed with cannabis use disorder: the moderating role of cognitive control.** *Addict Behav* 2020, **100**:106126 <http://dx.doi.org/10.1016/j.addbeh.2019.106126>.
24. O'Neill A, Bachi B, Bhattacharyya S: **Attentional bias towards cannabis cues in cannabis users: A systematic review and meta-analysis.** *Drug Alcohol Depend* 2020, **206**:107719 <http://dx.doi.org/10.1016/j.drugalcdep.2019.107719>.
- 24]O'Neill A, Bachi B, Bhattacharyya S: **Attentional bias towards cannabis cues in cannabis users: A systematic review and meta-analysis.** *Drug Alcohol Depend* 2020, **206**: 107719. <https://doi.org/10.1016/j.drugalcdep.2019.107719>. A systematic review and meta-analysis on attentional bias in cannabis users. While study heterogeneity is a problem and effect sizes are small, there is evidence that cannabis users show an attentional bias for cannabis words as well as pictures. Interestingly, effect sizes are bigger in studies where stimuli are presented for shorter periods of time, suggesting that attentional bias might be a predominantly fast and automatic process.
25. Cousijn J, Watson P, Koenders L, Vingerhoets WAM, Goudriaan AE, Wiers RW: **Cannabis dependence, cognitive control and attentional bias for cannabis words.** *Addict Behav* 2013, **38**:2825-2832 <http://dx.doi.org/10.1016/j.addbeh.2013.08.011>.
26. Ruglass LM, Shevorykin A, Dambreville N, Melara RD: **Neural and behavioral correlates of attentional bias to cannabis cues among adults with cannabis use disorders.** *Psychol Addict Behav* 2019, **33**:69-80 <http://dx.doi.org/10.1037/adb0000423>.
- 26]Ruglass LM, Shevorykin A, Dambreville N, Melara RD: **Neural and behavioral correlates of attentional bias to cannabis cues among adults with cannabis use disorders.** *Psychol Addict Behav* 2019, **33**: 6980. <https://doi.org/10.1037/adb0000423>. An EEG study (N = N40) investigating the neural mechanisms underlying attentional bias. Results show that individuals with a cannabis use disorder had poorer inhibition and selective attention than controls, and attentional bias for cannabis cues seems to be related to altered early perceptual processing.
27. Metrik J, Kahler CW, Reynolds B, McGeary JE, Monti PM, Haney M, de Wit H, Rohsenow DJ: **Balanced placebo design with marijuana: pharmacological and expectancy effects on impulsivity and risk taking.** *Psychopharmacology (Berl)* 2012, **223**:489-499 <http://dx.doi.org/10.1007/s00213-012-2740-y>.
28. McDonald J, Schleifer L, Richards JB, de Wit H: **Effects of THC on behavioral measures of impulsivity in humans.** *Neuropsychopharmacology* 2003, **28**:1356-1365 <http://dx.doi.org/10.1038/sj.npp.1300176>.
29. Broyd SJ, van Hell HH, Beale C, Yücel M, Solowij N: **Acute and chronic effects of cannabinoids on human cognition—a systematic review.** *Biol Psychiatry* 2016, **79**:557-567 <http://dx.doi.org/10.1016/j.biopsych.2015.12.002>.
30. Boggs DL, Cortes-Briones JA, Surti T, Luddy C, Ranganathan M, Cahill JD, Sewell AR, D'Souza DC, Skosnik PD: **The dose-dependent psychomotor effects of intravenous delta-9-tetrahydrocannabinol (Δ^9 -THC) in humans.** *J Psychopharmacol* 2018, **32**:1308-1318 <http://dx.doi.org/10.1177/0269881118799953>.
31. Kroon E, Kuhns L, Hoch E, Cousijn J: **Heavy cannabis use, dependence and the brain: a clinical perspective.** *Addiction* 2020, **115**:559-572 <http://dx.doi.org/10.1111/add.14776>.
32. Sofis MJ, Budney AJ, Stanger C, Knapp AA, Borodovsky JT: **Greater delay discounting and cannabis coping motives are associated with more frequent cannabis use in a large sample of adult cannabis users.** *Drug Alcohol Depend* 2020, **207**:107820 <http://dx.doi.org/10.1016/j.drugalcdep.2019.107820>.
33. Gilman JM, Calderon V, Curran MT, Evins AE: **Young adult cannabis users report greater propensity for risk-taking only in non-monetary domains.** *Drug Alcohol Depend* 2015, **147**:26-31 <http://dx.doi.org/10.1016/j.drugalcdep.2014.12.020>.
34. Fridberg DJ, Queller S, Ahn WY, Kim W, Bishara AJ, Busemeyer JR, Porrino L, Stout JC: **Cognitive mechanisms underlying risky decision-making in chronic cannabis users.** *J Math Psychol* 2010, **54**:28-38 <http://dx.doi.org/10.1016/j.jmp.2009.10.002>.
35. Wesley MJ, Hanlon CA, Porrino LJ: **Poor decision-making by chronic marijuana users is associated with decreased functional responsiveness to negative consequences.** *Psychiatry Res Neuroimaging* 2011, **191**:51-59 <http://dx.doi.org/10.1016/j.pscychresns.2010.10.002>.
36. Cuttler C, Spradlin A, Nusbaum AT, Whitney P, Hinson JM, McLaughlin RJ: **Joint effects of stress and chronic cannabis use on prospective memory.** *Psychopharmacology (Berl)* 2019, **236**:1973-1983 <http://dx.doi.org/10.1007/s00213-019-5184-9>.
37. Miranda R, Wemm SE, Treloar Padovano H, Carpenter RW, Emery NN, Gray JC, Mereish EH: **ives.** *Clin Psychol Sci* 2019, **7**:1094-1108 <http://dx.doi.org/10.1177/2167702619841976>.
- 37]Miranda R, Wemm SE, Treloar Padovano H, Carpenter RW, Emery NN, Gray JC, Mereish EH: **Weaker Memory Performance Exacerbates Stress-Induced Cannabis Craving in Youths Daily Lives.** *Clin Psychol Sci* 2019, **7**: 1094-1108. <https://doi.org/10.1177/2167702619841976>. A study assessing the association between working memory (WM), (cue-induced) craving and stress (N = N85), with results indicating that good WM might protect against increased craving under stressful circumstances.
38. Hindocha C, Freeman TP, Schafer G, Gardener C, Das RK, Morgan CJA, Curran HV: **Acute effects of delta-9-tetrahydrocannabinol, cannabidiol and their combination on facial emotion recognition: a randomised, double-blind, placebo-controlled study in cannabis users.** *Eur Neuropsychopharmacol* 2015, **25**:325-334 <http://dx.doi.org/10.1016/j.euroneuro.2014.11.014>.
39. Fusar-Poli P, Crippa JA, Bhattacharyya S, Borgwardt SJ, Allen P, Martin-Santos R, Seal M, Surguladze SA, O'Carroll C, Atakan Z et al.: **Distinct effects of Δ^9 -tetrahydrocannabinol and cannabidiol on neural activation during emotional processing.** *Arch Gen Psychiatry* 2009, **66**:95-105 <http://dx.doi.org/10.1001/archgenpsychiatry.2008.519>.
40. Bossong MG, van Hell HH, Jager G, Kahn RS, Ramsey NF, Jansma JM: **The endocannabinoid system and emotional processing: a pharmacological fMRI study with Δ^9 -tetrahydrocannabinol.** *Eur Neuropsychopharmacol* 2013, **23**:1687-1697 <http://dx.doi.org/10.1016/j.euroneuro.2013.06.009>.
41. Troup LJ, Andrzejewski JA, Torrence RD: **The effects of sex and residual cannabis use on emotion processing: An event-related potential study.** *Exp Clin Psychopharmacol* 2019, **27**:318-325 <http://dx.doi.org/10.1037/pha0000265>.
- 41]Troup LJ, Andrzejewski JA, Torrence RD: **The effects of sex and residual cannabis use on emotion processing: An event-related potential study.** *Exp Clin Psychopharmacol* 2109, **27**: 318325. <https://doi.org/10.1037/pha0000265>. An EEG study assessing the relationship between residual cannabis use, gender, and emotional processing (N = N144). Results show complex gender interactions that highlight the importance of assessing gender differences in the effects of cannabis on in particular emotion processing.
42. Figueiredo PR, Tolomeo S, Steele JD, Baldacchino A: **Neurocognitive consequences of chronic cannabis use: a systematic review and meta-analysis.** *Neurosci Biobehav Rev* 2020, **108**:358-369 <http://dx.doi.org/10.1016/j.neubiorev.2019.10.014>.
- 42]Figueiredo PR, Tolomeo S, Steele JD, Baldacchino A: **Neurocognitive consequences of chronic cannabis use: a systematic review and meta-analysis.** *Neurosci Biobehav Rev* 2020, **108**: 358369. <https://doi.org/10.1016/j.neubiorev.2019.10.014>.

- 10.1016/j.neubiorev.2019.10.014. A systematic review and meta-analysis assessing the evidence for the association between chronic cannabis use and neurocognitive functioning, finding small associations between chronic cannabis use and performance in a variety of cognitive domains, but not motor impulsivity. Problems with the heterogeneity in measures used to assess the same cognitive domain are discussed.
43. Fatima H, Howlett AC, Whitlow CT: **Reward, control & decision-making in cannabis use disorder: insights from functional MRI.** *Br J Radiol* 2019, **92**:20190165 <http://dx.doi.org/10.1259/bjr.20190165>.
 44. Karschner EL, Schwilke EW, Lowe RH, Darwin WD, Hering WI, Cadet JL, Huestis MA: **Implications of plasma 9-tetrahydrocannabinol, 11-hydroxy-THC, and 11-nor-9-carboxy-THC concentrations in chronic cannabis smokers.** *J Anal Toxicol* 2009, **33**:469-477 <http://dx.doi.org/10.1093/jat/33.8.469>.
 45. Sharma P, Murthy P, Bharath MMS: **Chemistry, metabolism, and toxicology of cannabis: clinical implications.** *Iran J Psychiatry* 2012, **7**:149-156.
 46. Bolla KI, Brown K, Eldreth D, Tate K, Cadet JL: **Dose-related neurocognitive effects of marijuana use.** *Neurology* 2002, **59**:1337-1343 <http://dx.doi.org/10.1212/01.WNL.0000031422.66442.49>.
 47. Casajuana Kögel C, Balcells-Olivero MM, López-Pelayo H, Miquel L, Teixidó L, Colom J, Nutt DJ, Rehm J, Gual A: **The standard joint unit.** *Drug Alcohol Depend* 2017, **176**:109-116 <http://dx.doi.org/10.1016/j.drugalcdep.2017.03.010>.
 48. Freeman TP, Lorenzetti V: **A proposal to standardize dose across all cannabis products and methods of administration.** *Addiction* 2019, **115**:1207-1216 <http://dx.doi.org/10.1111/add.14842>.
 - 48]Freeman TP, Lorenzetti V: **Standard THC units: a proposal to standardize dose across all cannabis products and methods of administration.** *Addiction* 2019, **115**: 1207-1216. <https://doi.org/10.1111/add.14842>. A recent effort introducing the Standard THC unit as a potential standardized measure of THC dose. Advantages include that it has the potential to be used across multiple products and administration methods. Disadvantages for research as well as implementation include the limitations that are put on analysing cannabis samples in jurisdictions where cannabis is not fully legal.
 49. UNODC: **Analysis of drug markets.** *World Drug Report 2018.* 2018:1-72.
 50. Littman R, Takács Á: **Do all inhibitions act alike? A study of go/no-go and stop-signal paradigms.** *PLoS One* 2017, **12**:1-20 <http://dx.doi.org/10.1371/journal.pone.0186774>.
 51. Cousijn J, Núñez AE, Filbey FM: **Time to acknowledge the mixed effects of cannabis on health: a summary and critical review of the NASEM 2017 report on the health effects of cannabis and cannabinoids.** *Addiction* 2018, **113**:958-966 <http://dx.doi.org/10.1111/add.14084>.
 52. Crean RD, Crane NA, Mason BJ: **An evidence-based review of acute and long-term effects of cannabis use on executive cognitive functions.** *J Addict Med* 2011, **5**:1-8 <http://dx.doi.org/10.1097/ADM.0b013e31820c23fa>.
 53. Zhang MWB, Ying J, Wing T, Song G, Fung DSS, Smith HE: **Cognitive biases in cannabis, opioid, and stimulant disorders: a systematic review.** *Front Psychiatry* 2018, **9**:376 <http://dx.doi.org/10.3389/fpsy.2018.00376>.