


## RESEARCH ARTICLE

# Safety and tolerability of natural and synthetic cannabinoids in adults aged over 50 years: A systematic review and meta-analysis

Latha Velayudhan<sup>1</sup>, Katie McGoohan<sup>1</sup>, Sagnik Bhattacharyya<sup>1,2\*</sup>

**1** Department of Old age Psychiatry, Division of Academic Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom, **2** Department of Psychosis Studies, Division of Academic Psychiatry, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

\* [Sagnik.2.bhattacharyya@kcl.ac.uk](mailto:Sagnik.2.bhattacharyya@kcl.ac.uk)



## Abstract

### Background

Cannabinoid-based medicines (CBMs) are being used widely in the elderly. However, their safety and tolerability in older adults remains unclear. We aimed to conduct a systematic review and meta-analysis of safety and tolerability of CBMs in adults of age  $\geq 50$  years.

### Methods and findings

A systematic search was performed using MEDLINE, PubMed, EMBASE, CINAHL PsychInfo, Cochrane Library, and ClinicalTrials.gov (1 January 1990 to 3 October 2020). Randomised clinical trials (RCTs) of CBMs in those with mean age of  $\geq 50$  years for all indications, evaluating the safety/tolerability of CBMs where adverse events have been quantified, were included. Study quality was assessed using the GRADE (Grading of Recommendations Assessment, Development, and Evaluation) criteria and Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines were followed. Two reviewers conducted all review stages independently. Where possible, data were pooled using random-effects meta-analysis. Effect sizes were calculated as incident rate ratio (IRR) for outcome data such as adverse events (AEs), serious AEs (SAEs), and death and risk ratio (RR) for withdrawal from study and reported separately for studies using tetrahydrocannabinol (THC), THC:cannabidiol (CBD) combination, and CBD. A total of 46 RCTs were identified as suitable for inclusion of which 31 (67%) were conducted in the United Kingdom and Europe. There were 6,216 patients (mean age  $58.6 \pm 7.5$  years; 51% male) included in the analysis, with 3,469 receiving CBMs. Compared with controls, delta-9-tetrahydrocannabinol (THC)-containing CBMs significantly increased the incidence of all-cause and treatment-related AEs: THC alone (IRR: 1.42 [95% CI, 1.12 to 1.78]) and (IRR: 1.60 [95% CI, 1.26 to 2.04]); THC:CBD combination (IRR: 1.58 [95% CI, 1.26 to 1.98]) and (IRR: 1.70 [95% CI, 1.24 to 2.33]), respectively. IRRs of SAEs and deaths were not significantly greater under CBMs containing THC

## OPEN ACCESS

**Citation:** Velayudhan L, McGoohan K, Bhattacharyya S (2021) Safety and tolerability of natural and synthetic cannabinoids in adults aged over 50 years: A systematic review and meta-analysis. *PLoS Med* 18(3): e1003524. <https://doi.org/10.1371/journal.pmed.1003524>

**Academic Editor:** Alexander C. Tsai, Massachusetts General Hospital, UNITED STATES

**Received:** October 7, 2020

**Accepted:** December 15, 2020

**Published:** March 29, 2021

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pmed.1003524>

**Copyright:** © 2021 Velayudhan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its [Supporting Information](#) files.

**Funding:** SB is supported by grants from the National Institute of Health Research (NIHR) Efficacy and Mechanism Evaluation scheme (UK) (grant number 16/126/53) and SB and LV are in receipt of funding from Parkinson's UK (grant number G-1901). The authors acknowledge support from the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

**Abbreviations:** AEs, adverse event; CBD, cannabidiol; CBM, cannabinoid-based medicine; GRADE, Grading of Recommendations Assessment, Development, and Evaluation; IRR, incident rate ratio; MedDRA, Medical Dictionary for Regulatory Activities; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-analyses; RCT, randomised clinical trial; RR, risk ratio; SAE, serious adverse event; SOC, system organ classes; THC, delta-9-tetrahydrocannabinol.

with or without CBD. THC:CBD combination (RR: 1.40 [95% CI, 1.08 to 1.80]) but not THC alone (RR: 1.18 [95% CI, 0.89 to 1.57]) significantly increased risk of AE-related withdrawals. CBD alone did not increase the incidence of all-cause AEs (IRR: 1.02 [95% CI, 0.90 to 1.16]) or other outcomes as per qualitative synthesis. AE-related withdrawals were significantly associated with THC dose in THC only [QM (df = 1) = 4.696,  $p = 0.03$ ] and THC:CBD combination treatment ([QM (df = 1) = 4.554,  $p = 0.033$ ]. THC-containing CBMs significantly increased incidence of dry mouth, dizziness/light-headedness, and somnolence/drowsiness. Study limitations include inability to fully exclude data from those <50 years of age in our primary analyses as well as limitations related to weaknesses in the included trials particularly incomplete reporting of outcomes and heterogeneity in included studies.

## Conclusions

This pooled analysis, using data from RCTs with mean participant age  $\geq 50$  years, suggests that although THC-containing CBMs are associated with side effects, CBMs in general are safe and acceptable in older adults. However, THC:CBD combinations may be less acceptable in the dose ranges used and their tolerability may be different in adults over 65 or 75 years of age.

## Author summary

### Why was this study done?

- Use of cannabinoid-based medicines (CBMs) has been growing steadily in recent years, including in the elderly. However, their safety and tolerability in older adults remains unclear.
- With increasing interest in the use of CBMs in older people and growing unlicensed use, there is a particular need to examine their safety and tolerability in older adults.
- We analysed data on safety and tolerability from previously published double-blind, randomised controlled trials (RCT) using delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD), the common constituents of most CBMs, alone or in combination, to examine their effect on older adults.

### What did the researchers do and find?

- We pooled data from 46 published RCTs (with information from 6,216 patients; with mean participant age  $\geq 50$  years) on adverse events, serious adverse events or death, and withdrawal from study. We also examined the relationship between the dose of THC used in THC-containing CBMs and the incidence of adverse consequences in older adults.

- Our results suggest that compared with the control condition, treatment with THC-containing CBMs was associated on average with significantly greater incidence of all-cause and treatment-related adverse events.
- There was no significant increase in the incidence of serious adverse events or death with any CBMs. The risk of withdrawal from study was increased only in those receiving THC:CBD combination treatment, and this was related to THC dose.

### What do these findings mean?

- These findings suggest that CBMs in general are safe and acceptable in older adults.
- Our findings that THC-containing CBMs are associated with side effects and that THC:CBD combinations may be less acceptable at the dose ranges typically used in RCTs is critical to prescribing in older people.

## Introduction

The cannabis plant (*Cannabis sativa* L.) has been used worldwide both for recreational and medicinal purposes for thousands of years. With a fast-growing aging population, its medicinal use has also caught up and is growing in the elderly [1–3].

Among the cannabinoids found in the cannabis plant, delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) are the most well characterised and often considered for medicinal purposes. THC can cause intoxication [4,5] and has antiemetic, analgesic, and potentially neuroprotective and anti-inflammatory effects. On the other hand, CBD is nonintoxicating [5,6] with antiepileptic and potentially also anti-inflammatory, neuroprotective, antioxidant, and antipsychotic effects [7–9]. While several trials have used these cannabinoids for a wide range of diseases and indications, a majority of these have investigated younger people [10,11]. However, age-related pharmacodynamic and pharmacokinetic changes as well as higher prevalence of comorbidities and polypharmacy in the elderly mean that they may have a different profile of safety and tolerability to cannabinoids [12,13] compared to younger people, as is well known with other groups of medications especially those used for disorders of the central nervous system [14]. Both THC and CBD, the common constituents of most cannabinoid-based medicines in current use have prominent effects on brain function and cognition [15]. Therefore, evidence of safety and tolerability of cannabinoid-based medicines (CBM) established in studies in younger adults cannot be directly extrapolated to the older adults. Although a number of recent reviews and meta-analyses [12,16,17] have summarized the safety and tolerability profile of CBMs, they have all pooled data from studies investigating across the age spectrum, making it difficult to draw age-specific inferences. With increasing interest in their use in disorders typically affecting older people [18–20] and growing unlicensed use [21], there is a particular need to investigate the safety and tolerability of CBMs in older people. This is also relevant, as there is a widely held view that many of the naturally derived cannabinoids are generally safe as they have been around and used for a long time.

Here, we have addressed this by investigating the safety and tolerability of CBMs in people over 50 years of age through systematically reviewing all double-blind, randomised controlled

trials (RCTs) using CBMs that focused on people with mean age of 50 years and over to conduct a meta-analysis. As there is a larger evidence base of studies with mean age of participants  $\geq 50$  years than the more limited set of studies that have exclusively focused on people over 50 years and even less on people over 65 or 75 years, we have focused on studies with mean age of participants  $\geq 50$  years and complemented these results with additional analyses restricted to studies that have exclusively focused on people over 50 years and even less on people over 65. Existing meta-analytic investigations [16,17] have generally considered all CBMs together, irrespective of whether they included THC, CBD, or THC:CBD in combination. However, THC can cause intoxication and may induce anxiety and transient psychotomimetic effects [5], especially at higher doses and in vulnerable individuals, while CBD does not cause intoxication when directly compared in the same individuals [5] and may potentially ameliorate anxiety and psychosis [9,22–24]. Further, there is growing evidence that THC and CBD may have opposing acute effects on autonomic arousal and brain [15] and cardiovascular function [25,26], and CBD may mitigate some of the harmful effects of THC on cognition and behaviour [15,27,28], consistent with their opposing effects on some of their molecular targets [4]. This suggests that THC and CBD may have distinct tolerability profiles, with the possibility that certain side effects may be noticeable in those taking formulations containing only THC but not in those taking formulations containing only CBD while adverse effects may even be mitigated in those taking THC and CBD in combination. This underscores the importance of examining their safety and tolerability separately. Therefore, we have addressed this issue by separately investigating the effects of THC, CBD, or THC:CBD in combination.

We hypothesized that compared to control treatments, all 3 categories of CBMs will be associated with: (i) a greater incidence of adverse events (AEs); (ii) no greater incidence of serious adverse events (SAEs) or death; and (iii) no greater risk of withdrawal from study. Further, we hypothesized a direct relationship between the dose of THC used in THC-containing CBMs and the incidence of adverse consequences in older adults.

## Methods

### Data sources and searches

The review was undertaken according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guidelines [29] (see [S1 PRISMA checklist](#)). The study protocol was preregistered with the International Prospective Register of Systematic Reviews (PROSPERO) (CRD42019148869). Ethics approval was not required for this systematic review and meta-analysis.

A detailed description of the bibliographic search strategy is presented in Methods in [S1 Text](#). We identified studies published from 1 January 1990 up to 31 October 2020, from several electronic databases. Studies were independently assessed by 2 researchers and disagreements resolved through consensus or discussions with a third researcher.

### Study selection

Studies were included if (1) published from 1990 onwards; (2) included older adults (defined as mean age  $\geq 50$  years) or reported a distinct subgroup of older adults and provided separate results for this subgroup; and (3) provided data on the safety and tolerability of medical cannabinoids administered by any route, at any dose, for any duration and for any indication. Studies were excluded if they (1) included exclusively younger participants (mean age  $< 50$  years); (2) studied effects of cannabinoids for recreational purposes or failed to provide the dosage of cannabinoids; and (3) were not reported in English language. Here, we focus on results from RCTs.



## Data extraction and quality assessment

All relevant available data for examination of the safety and tolerability of different CBMs (THC:CBD combination or THC or CBD alone) were collected from eligible studies, complemented with information from [ClinicalTrials.gov](https://clinicaltrials.gov) and author responses.

Data were extracted for study design, participant characteristics, indication, dosage and duration of intervention, all-cause and treatment-related AEs and SAEs, and AE-related withdrawals and deaths.

AEs and SAEs were coded according to the Medical Dictionary for Regulatory Activities (MedDRA) “system organ classes” (SOC). Data were also extracted for the top 5 (as reported by each study) AEs for each SOC, where available. Data extraction and coding was verified by a medically qualified researcher and discrepancies resolved following discussions with senior researcher. The disease conditions investigated were classified into broader subgroups for analysis purpose.

Overall, quality of evidence was assessed using recommended criteria [30] and summarised to reflect confidence in estimates [31].

## Data synthesis and analysis

Total exposure to active intervention in person-years was estimated by first calculating this for each individual study by multiplying the number of participants in the active intervention arm with the duration of treatment for that arm for each study and then adding up these study-specific values for all studies under each broad category (THC, THC:CBD, or CBD) of intervention investigated here. Mean exposure in person-years for each category (THC, THC:CBD, or CBD) of intervention was estimated by calculating the arithmetic mean from study-specific estimates obtained as above for each intervention category. Pooled mean ages of participants for each group of studies and treatment arms were estimated by calculating the arithmetic mean of study specific mean age as reported by individual studies for each intervention arm. Other pooled estimates (median and interquartile range) for summary study characteristics, such as duration of study in weeks, participants analysed or included, or duration of treatment for each treatment arm, were calculated by estimating them from the total number (e.g., for variables such as number of participants included or analysed; duration of study or treatment) or the mean estimate (e.g., for variables such as mean age) as reported in individual studies for each set subgroups of RCTs.


Pooled effect-sizes were estimated if there were 2 or more RCTs within each group or subgroup under the random-effects model using the restricted maximum-likelihood estimator because of anticipated heterogeneity. For each broad category of intervention, analyses combined both parallel-arm and crossover RCTs, with the latter treated as parallel-arm design [32] for pooled analyses. But we also report results by RCT design for each intervention. We estimated incident rate ratio (IRR) for outcome data such as AEs, SAEs, and death and risk ratio (RR) for withdrawal from study. Studies with more than 1 active treatment arm were treated as independent studies. In studies with more than 1 active treatment arm compared against a single control group, we also report meta-analysis of dependent effect-sizes with robust variance estimation [33–35]. For the purpose of reporting throughout the manuscript, results are reported for analyses treating all studies as independent, while corresponding dependent meta-analyses are reported in Results in [S1 Text](#) and signposted in the main text as appropriate. We investigated heterogeneity using forest plots and the  $I^2$  statistic and publication bias using Egger regression test [36] and the “Trim and fill” method [37]. For the analysis of AEs, data for all conditions were combined. We also examined whether estimates varied according to treatment, design, clinical condition, and dose of study drug using meta-regression.

Our primary analysis includes the results of all studies where the mean age of study participants was  $\geq 50$  years. As many participants in these included studies were  $< 50$  years of age, we also carried out separate analyses restricted to studies where all participants were  $\geq 50$  years of age and also where all participants were  $\geq 65$  years of age. These analyses were carried out where there were at least 2 studies with analysable data. Statistical analyses were performed using the *metafor* package in R (version 3.6.3) [38]. For meta-analysis of dependent effect-sizes with robust variance estimation, we also used the *clubSandwich* package (<https://github.com/jepusto/clubSandwich>) along with *metafor* in R.

## Results

### Data selection

[Fig 1](#) (PRISMA flow chart) summarizes the study selection procedure. Main characteristics and outcome measures of each study are included in [Tables 1–3](#); additional details regarding



**Fig 1. Study disposition.**

<https://doi.org/10.1371/journal.pmed.1003524.g001>

Table 1. Characteristics of included randomised controlled trials of THC in older adults ( $N = 30$ ).

Study ID (country)	Study Design (RCT)	THC: Sample included/analysed N Mean age (SD), Male %	Comparator: Sample included/analysed N Mean age (SD), Male %	Age cut-off for enrolment	Indication	THC classification	Comparator	THC treatment duration, weeks	Calculated daily average THC dose	Overall GRADE rating for study
*** Ahmed et al. 2014 (the Netherlands)	Crossover	12/11 72.00 (5), 50	12/11 72.00 (5), 50	≥65 years	Healthy older participants	Namisol	Placebo	.4	6.5 mg	Moderate
Ahmed et al. 2015 (the Netherlands)	Crossover	10/10 77.30 (5.6), 70	10/10 77.30 (5.6), 70	≥18 years	Dementia	Namisol	Placebo	2.6	3 mg	Moderate
Brisbois et al. 2011 (Canada)	Parallel-arm	24/11 67.00 (10.9), 64	22/10 65.50 (8), 50	Adult	Cancer patients with chemosensory alterations	Dronabinol	Placebo	2.6	7.5 mg	Low
Carley et al. 2018 (USA) <sup>‡</sup>	Parallel-arm	21/21 52.70 (7.7), 76	25/25 58.80 (6.1), 72	21–65 years	Obstructive sleep apnoea	Dronabinol	Placebo	6.0	2.5 mg	Low
Carley et al. 2018 (USA) <sup>‡</sup>	Parallel-arm	27/27 54.70 (7), 67	25/25 58.80 (6.1), 72	21–65 years	Obstructive sleep apnoea	Dronabinol	Placebo	6.0	10 mg	Low
Curtis et al. 2009 (UK)	Crossover	44/37 52.00 (9.5), 50	44/37 52.00 (9.5), 50	≥18 years	Huntington disease	Nabilone	Placebo	5.0	2 mg	Low
De Vries et al. 2016 (the Netherlands)	Crossover	25/24 52.00 (NR), 62	25/24 52.00 (NR), 62	>18 years	Chronic pancreatitis	Namisol	Diazepam	.1	8 mg	Moderate
** Herrmann et al. 2019 (Canada)	Crossover	39/38 87.00 (10), 77	39/38 87.00 (10), 77	≥55 years	Alzheimer disease	Nabilone	Placebo	6.0	1.6 mg	Moderate
Jatoi et al. 2002 (USA)	Parallel-arm	152/152 67.00 (10), 66	159/159 65.00 (11), 65	≥18 years	Cancer-related anorexia	Dronabinol	Megestrol acetate	8.1	5 mg	Low
Johnson et al. 2010 (UK) <sup>§</sup>	Parallel-arm	58/58 61.30 (12.5), 52	59/59 60.10 (12.3), 54	NR	Patients with cancer-related pain	THC extract spray	Placebo	2.0	23 mg	Moderate
Lane et al. 1991 (USA)	Parallel-arm	21/21 47.0 (20–68) <sup>*,†</sup> 48	21/21 49.0 (22–64) <sup>*,†</sup> 48	18–69 years	Chemotherapy-induced nausea and vomiting	Dronabinol	Prochlorperazine	.9	40 mg	Low
Meiri et al. 2007 (USA)	Parallel-arm	17/17 61.60 (14.2), 53	14/14 57.20 (8.6), 38	≥18 years	Chemotherapy-induced nausea and vomiting	Dronabinol	Placebo	.7	20 mg	Low
Peball et al. 2020 (Austria)	Parallel-arm	19/19 65.4 (7.94), 53	19/19 64.0 (8.04), 74	≥30 years	Parkinson disease	Nabilone	Placebo	4.0	0.75 mg	Moderate
Sieradzan et al. 2001 (UK)	Crossover	9/9 59.00 (NR), 44	9/9 59.00 (NR), 44	NR	Parkinson disease	Nabilone	Placebo	.1	2 mg	Very low
Strasser et al. 2006 (Germany) <sup>§</sup>	Parallel-arm	100/100 60.00 (12), 54	48/48 62.00 (10), 52	Adult	Cancer-related anorexia	THC	Placebo	6.0	5 mg	Low
Svensen et al. 2004 (Denmark)	Crossover	24/24 50.0 (NR), 42	24/24 50.0 (NR), 42	18–55 years	Multiple sclerosis	Dronabinol	Placebo	3.0	10 mg	Moderate
Tomida et al. 2006 (UK) <sup>§</sup>	Crossover	6/6 55.30 (5), 100	6/6 55.30 (5), 100	NR	Intraocular pressure	THC extract spray	Placebo	.1	5 mg	Low
Toth et al. 2012 (Canada)	Parallel-arm	13/13 60.80 (15.3), 38	13/13 61.60 (14.6), 69	18–80	Diabetic peripheral neuropathic pain	Nabilone	Placebo	5.0	4 mg	Low

(Continued)

Table 1. (Continued)

Study ID (country)	Study Design (RCT)	THC: Sample included/analysed N Mean age (SD), Male %	Comparator: Sample included/analysed N Mean age (SD), Male %	Age cut-off for enrolment	Indication	THC classification	Comparator	THC treatment duration, weeks	Calculated daily average THC dose	Overall GRADE rating for study
Van Amerongen et al. 2017, 2 (the Netherlands) <sup>  </sup>	Crossover	24/24 54.30 (8.9), 33	24/24 54.30 (8.9), 33	≥18 years	Multiple sclerosis	THC	Placebo	.1	16 mg	Moderate
Van Amerongen et al. 2017, 1 (the Netherlands) <sup>  </sup>	Parallel-arm	12/12 57.30 (9), 33	12/12 51.40 (8), 33	≥18 years	Multiple sclerosis	THC	Placebo	4.0	28.5 mg	Moderate
Van den Elsen et al. 2015, 1 (the Netherlands)	Parallel-arm	24/24 79.00 (8), 46	26/26 78.00 (7), 54	≥40 years	Dementia	Namisol	Placebo	3.0	4.5 mg	Moderate
Van den Elsen et al. 2015, 2 (the Netherlands)	Crossover	22/22 76.40 (5.3), 68	22/22 76.40 (5.3), 68	≥18 years	Dementia	Namisol	Placebo	2.6	3 mg	Moderate
***Volicer et al. 1997 (USA)	Crossover	15/12 72.70 (4.9), 92	15/12 72.70 (4.9), 92	NR	Alzheimer disease	Dronabinol	Placebo	6.0	5 mg	Very low
***Walther et al. 2011 (Switzerland)	Crossover	2/2 78.00 (NR), 100	2/2 78.00 (NR), 100	NR	Alzheimer disease	Dronabinol	Placebo	2.0	2.5 mg	Very low
Ware et al. 2010 (Canada)	Crossover	32/32 50.00 (11.2), 16	32/32 50.00 (11.2), 16	≥18 years	Fibromyalgia	Nabilone	Amitriptyline	2.0	1 mg	Moderate
Weber et al. 2010 (Switzerland)	Crossover	27/22 57.00 (12), 74	27/22 57.00 (12), 74	Adult	Amyotrophic lateral sclerosis patients with cramps	Dronabinol	Placebo	2.0	10 mg	Moderate
Zadikoff et al. 2011 (Canada)	Crossover	9/9 60.00 (7), 0	9/9 60.00 (7), 0	18–75	Cervical dystonia	Dronabinol	Placebo	3.0	15 mg	Low
Zajicek et al. 2003 (UK) <sup>§</sup>	Parallel-arm	216/206 50.00 (8.2), 31	222/213 51.00 (7.6), 37	18–64	Multiple sclerosis	Dronabinol	Placebo	14.0	25mg	Moderate
Zajicek et al. 2005 (UK) <sup>§</sup>	Parallel-arm	125/125 50.00 (8.2), 31	120/120 51.00 (7.6), 37	18–64	Multiple sclerosis	Dronabinol	Placebo	52.0	25 mg	Moderate
Zajicek et al. 2013 (UK)	Parallel-arm	332/329 52.30 (7.6), 40	166/164 52.00 (8.2), 41	18–65	Multiple sclerosis	Dronabinol	Placebo	160.0	28 mg	Moderate

\*\*\*Studies recruited participants ≥65 years.

‡Article included more than 1 dose level.

§Article included more than 1 cannabinoid intervention.

\*Median age (range).

†Included as median age for whole study population was ≥50.

||Article included the results of multiple trials.

GRADE, Grading of Recommendations Assessment, Development, and Evaluation; NR, not recorded; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.t001>

studies are presented in Results in [S1 Text](#) and Tables A and B in [S1 Text](#). A total of 60 comparisons of CBM and control intervention using RCT design (hereafter called RCTs) ( $n = 6,216$  participants; 1933.47 person-years of cannabinoid exposure) from 46 published articles

Table 2. Characteristics of included randomised controlled trials of THC:CBD combination in older adults (N = 26).

Study ID (country)	Study Design	CBD/THC: Sample included/analysed N Mean age (SD), Male %	Comparator: Sample included/analysed N Mean age (SD), Male %	Age cut-off for enrolment	Indication	CBD/THC classification	Comparator	CBD/THC treatment duration, weeks	Calculated daily average CBD/THC dose	GRADE rating
Blake et al. 2006 (UK)	Parallel-arm	31/31 60.9 (10.6), 26	27/27 64.9 (8.5), 15	NR	Rheumatoid arthritis	THC:CBD spray	Placebo	5.0	14.6 mg THC: 13.5 mg CBD	Low
**Carroll et al. 2004 (UK)	Crossover	19/17 67.0 (NR) 63	19/17 67.0 (NR) 63	18–78 years	Levodopa-induced dyskinesia in Parkinson disease	Cannabis extract	Placebo	4.0	10.2 mg THC: 5.1 mg CBD	Moderate
Duran et al. 2010 (Spain)	Parallel-arm	7/7 50 (41–70)* 0	9/9 50 (34–76)* 11	>18 years	Chemotherapy-induced nausea and vomiting	THC:CBD spray	Placebo	.6	13 mg THC: 12 mg CBD	Moderate
Fallon et al. 2017, 1 (Multicentre) <sup>  </sup>	Parallel-arm (withdrawal study)	103/103 61.4 (10.9), 61	103/103 61.6 (11.8), 53	≥18 years	Advanced cancer patients with pain	THC:CBD spray	Placebo	5.0	17.6 mg THC: 16.3 mg CBD	Moderate
Fallon et al. 2017, 2 (Multicentre) <sup>  </sup>	Parallel-arm	200/199 60.0 (11), 53	199/198 59.6 (11), 49	≥18 years	Advanced cancer patients with pain	THC:CBD spray	Placebo	5.0	17 mg THC: 15.8 mg CBD	Moderate
Jadoon et al. 2016, 1 (UK) <sup>‡,§</sup>	Parallel-arm	11/11 59.0 (8.8), 55	14/14 59.0 (7.7), 50	≥18 years	Type 2 diabetes	CBD/THCV	Placebo	13.0	10 mg THC: 10 mg CBD	Moderate
Jadoon et al. 2016, 2 (UK) <sup>‡,§</sup>	Parallel-arm	12/12 58.0 (8.1), 75	14/14 59.0 (7.7), 50	≥18 years	Type 2 diabetes	CBD/THCV	Placebo	13.0	10 mg THC: 200 mg CBD	Moderate
Johnson et al. 2010, (UK) <sup>§</sup>	Parallel-arm	60/60 59.4 (12.1), 55	59/59 60.1 (12.3), 54	NR	Patients with cancer-related pain	THC:CBD spray	Placebo	2.0	25 mg THC: 23 mg CBD	Moderate
Litchman et al. 2018, (Multicentre)	Parallel-arm	199/199 59.2 (12), 56	198/198 60.7 (11.1), 52	≥18 years	Advanced cancer patients with pain	THC:CBD spray	Placebo	5.0	17.3 mg THC: 16 mg CBD	Moderate
Lynch et al. 2014 (USA)	Crossover	18/16 56.0 (10.8), 17	18/16 56.0 (10.8), 17	NR	Chemotherapy-induced neuropathic pain	THC:CBD spray	Placebo	6.0	21.6 mg THC: 20 mg CBD	Low
Markova et al. 2019 (Czech Republic)	Parallel-arm	53/53 51.3 (10.2) 30	53/53 51.3 (10.2) 30	≥18 years	Multiple sclerosis	THC:CBD spray	Placebo	12.0	19.7 mg THC: 18.3 mg CBD	Low
Notcutt et al. 2012 (UK)	Parallel-arm (withdrawal study)	18/18 59.7 (9) 50	18/18 54.4 (10.4) 33	NR	Multiple sclerosis	THC:CBD spray	Placebo	4.0	20.8 mg THC: 19.3 mg CBD	Very low
Nurmikko et al. 2007 (UK)	Parallel-arm	63/63 52.4 (15.8), 44	62/62 54.3 (15.2), 37	≥18 years	Neuropathic pain	THC:CBD spray	Placebo	5.0	THC 29.7 mg: CBD 27.5 mg	High
***Pickering et al. 2011, 1 (UK) <sup>§</sup>	Crossover	5/4 67.0 (NR), 50	5/4 67.0 (NR), 50	40–74 years	COPD	THC:CBD spray	Placebo	.1	4.7 mg THC: 4.4 mg CBD	Low
**Pickering et al. 2011, 2 (UK) <sup>§</sup>	Crossover	6/5 58.0 (NR), 80	6/5 58.0 (NR), 80	40–75 years	Healthy controls	THC:CBD spray	Placebo	.1	10.3 mg THC: 9.5 mg CBD	Low
Portenoy et al. 2012, 1 (Multicentre) <sup>‡</sup>	Parallel-arm	91/91 59.0 (12.3), 49	91/91 56.0 (12.2), 48	NR	Cancer patients with chronic pain	THC:CBD spray	Placebo	5.0	10.8 mg THC: 10 mg CBD	Moderate

(Continued)



Table 2. (Continued)

Study ID (country)	Study Design	CBD/THC: Sample included/analysed N Mean age (SD), Male %	Comparator: Sample included/analysed N Mean age (SD), Male %	Age cut-off for enrolment	Indication	CBD/THC classification	Comparator	CBD/THC treatment duration, weeks	Calculated daily average CBD/THC dose	GRADE rating
Portenoy et al. 2012, 2 (Multicentre) <sup>‡</sup>	Parallel-arm	88/87 59.0 (13.1), 56	91/91 56.0 (12.2), 48	NR	Cancer patients with chronic pain	THC:CBD spray	Placebo	5.0	27 mg THC: 25 mg CBD	Moderate
Portenoy et al. 2012, 3 (Multicentre) <sup>‡</sup>	Parallel-arm	90/90 58.0 (11.2), 53	91/91 56.0 (12.2), 48	NR	Cancer patients with chronic pain	THC:CBD spray	Placebo	5.0	43.2 mg THC: 40 mg CBD	Moderate
Riva et al. 2019 (Italy)	Parallel-arm	30/29 58.4 (10.6) 62	30/30 57.2 (13.8) 53	18–80 years	Motor neurone disease	THC:CBD spray	Placebo	6.0	21.6 mg THC: 20.0 mg CBD	High
Serpell et al. 2014 (UK)	Parallel-arm	128/128 57.6 (14.4), 34	118/118 57.0 (14.1), 45	≥18 years	Neuropathic pain	THC:CBD spray	Placebo	14.0	24 mg THC: 22 mg CBD	Moderate
Strasser et al. 2006 (Germany) <sup>§</sup>	Parallel-arm	95/95 61.0 (12), 56	48/48 62.0 (10), 52	Adult	Cancer-related anorexia	Cannabis extract	Placebo	6.0	5 mg THC: 2 mg CBD	Moderate
Vaney et al. 2004 (Switzerland)	Crossover	57/50 55.0 (10), 49	57/50 55.0 (10), 49	Adult	Multiple sclerosis	Cannabis extract	Placebo	2.0	27.5 mg THC: 9.9 mg CBD	Low
Wade et al. 2004 (UK)	Parallel-arm	80/80 51.0 (9.4), 41	80/80 50.0 (9.3), 35	NR	Multiple sclerosis	THC:CBD spray	Placebo	6.0	40.5 mg THC: 37.5 mg CBD	Moderate
Zajicek et al. 2003 (UK) <sup>¶</sup>	Parallel-arm	219/211 51.0 (7.6), 36	222/213 51.0 (7.6), 37	18–64 years	Multiple sclerosis	Cannabis extract	Placebo	14.0	25 mg THC: 12.5 mg CBD	Moderate
Zajicek et al. 2005 (UK) <sup>¶</sup>	Parallel-arm	138/138 51.0 (7.6), 36	120/120 51.0 (7.6), 37	18–64 years	Multiple sclerosis	Cannabis extract	Placebo	52.0	25 mg THC: 12.5 mg CBD	Moderate
Zajicek et al. 2012 (UK)	Parallel-arm	144/143 51.9 (7.7), 39	135/134 52.0 (7.9), 35	18–64 years	Multiple sclerosis	Cannabis extract	Placebo	12.0	25 mg THC: 12.5 mg CBD	Moderate

\*\*Studies recruited participants ≥50years.

\*Median age (range).

||Article included the results of multiple trials.

‡Article included more than 1 dose level.

§Article included more than 1 cannabinoid intervention.

¶¶Studies recruited participants ≥65 years.

¶Article included multiple study groups/indications.

CBD, cannabidiol; COPD, chronic obstructive pulmonary disease; GRADE, Grading of Recommendations Assessment, Development, and Evaluation; NR, not recorded; THC, delta-9-tetrahydrocannabinol; THCV, tetrahydrocannabivarin.

<https://doi.org/10.1371/journal.pmed.1003524.t002>

were included (Fig 1). Of these, 4 RCTs recruited participants over age ≥65 years ( $n = 68$ ; mean age, 72.4 (SD ± 4.5)), of which one was ≥75 years (Tables 1 and 2) [39–42].

The formulations used in THC studies were (numbers within brackets indicating the number of RCTs where each formulation was used): nabilone (6), dronabinol (marinol) (14), THC (3), THC extract spray (2), and Namisol (5). The combination THC:CBD trials used THC:

Table 3. Characteristics of randomised controlled trials of CBD in older adults ( $N = 4$ ).

Study ID (country)	Study Design	CBD: Sample included/analysed N Mean age (SD) Male %	Comparator: Sample included/analysed N Mean age (SD) Male %	Age cut-off for enrolment	Indication	Active treatment	Comparator	CBD treatment duration, weeks	Calculated daily average CBD dose	GRADE rating
Consroe et al. 1991 USA	Crossover	18/15 47.8 <sup>2</sup> (15.3), 53	18/15 47.8 <sup>2</sup> (15.3), 53	NR	Huntington disease	CBD	Placebo	6	700 mg	Low
Jadoon et al. 2016 <sup>§</sup> UK	Parallel-arm	13/13 56.8 (9.9), 77	14/14 59.0 (9.4) 68	≥18 years	Type 2 diabetes	CBD	Placebo	13	200 mg	Moderate
Tomida et al. 2006 <sup>‡,§</sup> UK	Crossover	6/6 55.3 (5.0), 100	6/6 55.3 (5.0), 100	NR	Intraocular pressure	CBD	Placebo	0.1	20 mg	Low
Tomida et al. 2006 <sup>‡,§</sup> UK	Crossover	6/6 55.3 (5.0), 100	6/6 55.3 (5.0), 100	NR	Intraocular pressure	CBD	Placebo	0.1	40 mg	Low

§Article included more than 1 cannabinoid intervention.

‡Article included more than 1 dose level.

CBD, cannabidiol; GRADE, Grading of Recommendations Assessment, Development, and Evaluation; NR, not recorded.

<https://doi.org/10.1371/journal.pmed.1003524.t003>

CBD spray (18), cannabis extract (6), and CBD/THCV (2). The CBD studies used CBD preparations only.


The disease conditions investigated were classified into broader subgroups for analysis purpose as neurodegenerative (Alzheimer disease, Parkinson disease, Huntington disease, amyotrophic lateral sclerosis), multiple sclerosis, motor neurone disease, pain (neuropathic pain), cancer (cancer or chemotherapy-related anorexia, pain, or nausea/vomiting), and others (type 2 diabetes mellitus, chronic obstructive pulmonary disease, fibromyalgia, raised intraocular pressure, cervical dystonia, healthy, pancreatitis, obstructive sleep apnoea, and Levodopa-induced dyskinesia in Parkinson disease).

Figs 2–14 show the forest plots and summary results of the meta-analyses stratified according to study design, for all-cause and treatment-related AEs and SAEs, withdrawals, deaths, for studies using THC, THC:CBD combination, and CBD, respectively.

## THC studies

A total of 30 RCTs (15 crossover and 15 parallel-arm) from 28 articles [39–41,43–67] (see Results in S1 Text and Table A in S1 Text, for additional details), reported on 1,461 patients on active [analysed 1,417; Total person-years of THC exposure: 1252.83 person-years; Mean person-years of THC exposure (mean ± SD): 41.76 ± 184.28 person-years] and 1,251 (analysed 1,210) patients on control intervention, ranging from 50 to 87 years in mean age (males: 0% to 100%). All except 4 studies used placebo control [43,45,54,62].

Pooled IRRs for all-cause ( $k = 21$ ) and treatment-related AEs ( $k = 9$ ) from all RCTs were 1.42 (95% CI, 1.12 to 1.79) and 1.60 (95% CI, 1.26 to 2.04), respectively. Pooled IRRs for all-cause ( $k = 27$ ) and treatment-related ( $k = 23$ ) SAEs from all RCTs were 1.08 (95% CI, 0.80 to 1.46) and 1.23 (95% CI, 0.56 to 2.69), respectively. Pooled RR for AE-related withdrawals




**Fig 2. Forest plot of all-cause adverse events: THC studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g002>

(*k* = 27) and IRR for all deaths (*k* = 30) from all RCTs were 1.18 (95% CI, 0.89 to 1.57) and 1.09 (95% CI, 0.75 to 1.59), respectively. Neither Egger test nor “Trim and fill” method indicated publication or other selection bias for any of the outcomes except for SAEs (Results in [S1 Text](#) and Fig A in [S1 Text](#)). For all-cause SAEs, while Egger test was nonsignificant, Trim-fill method identified 10 missing studies. The estimated effect of treatment on IRR for all-cause SAEs, which was not significant previously, became significant after inclusion of potentially missing studies identified by the Trim-fill method (1.46, 95% CI, 1.09 to 1.95, *p* = 0.01, *k* = 35). For treatment-related SAEs, Egger test indicated significant publication bias and Trim-fill method identified 5 missing studies, though they did not change the direction or significance of effect-size on inclusion. Where there was nonindependence of outcome data used in analyses, results of dependent meta-analyses were consistent with the results of independent meta-analyses (Results in [S1 Text](#)).

**Effect of moderators.** Meta-regression analyses indicate that there was a trend-level effect [QM (df = 4) = 9.986, *p* = 0.084] of clinical condition on estimated effect of THC treatment on all-cause AEs, which seemed to be mainly related to a significantly lower estimated effect in RCTs investigating neurodegenerative disorder (regression coefficient = 0.905; *p* = 0.006)



**Fig 3. Forest plot of treatment-related adverse events: THC studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.


<https://doi.org/10.1371/journal.pmed.1003524.g003>

patients compared to other conditions. Except that, moderators such as study design or type of intervention did not significantly influence estimated effects of THC treatment on any of the outcomes assessed.

**Effect of dose.** Meta-regression analyses also indicated that there was a significant effect of daily THC dose on all-cause AEs [QM (df = 1) = 5.024, p = 0.025] as well as on AE-related withdrawals [QM (df = 1) = 4.696, p = 0.03] for all RCTs indicating that higher the dose of THC the higher was the risk of all-cause AEs and risk of withdrawal (regression coefficient = -0.905; p = 0.006) from study in THC-treated patients compared to control treatment (Fig B in S1 Text). There was no significant association of daily THC dose with any of the other estimates (SAEs and deaths).

**Common side effects.** Pooled IRRs of the most commonly reported AEs (Table C in S1 Text) suggested significantly higher incidence rate of dry mouth, dizziness/light-headedness, mobility/balance/coordination difficulties, somnolence/drowsiness, euphoria, and male impotence in active compared to control arms.

**Analysis of studies where all participants were ≥50 years.** Restricting the meta-analysis to the 4 studies [39–41,65] that recruited participants with ≥50 years of age (total n = 136; analysed n = 126), results were broadly comparable to the primary analysis in the pattern of




**Fig 4. Forest plot of all-cause serious adverse events: THC studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g004>

findings, though effect on all-cause AEs was no longer significant. The pooled IRRs for all-cause AEs (k = 3) were 2.13 (95% CI, 0.46 to 9.97) and treatment-related AEs (k = 3) was 2.80 (95% CI, 1.09 to 7.21), respectively. Pooled IRRs for all-cause (k = 4) and treatment-related SAEs (k = 2) were 1.20 (95% CI, 0.39 to 3.65) and 0.50 (95% CI, 0.03 to 7.99), respectively. Pooled RR for AE-related withdrawals (k = 4) and IRR for all deaths (k = 4) were 1.11 (95% CI, 0.49 to 2.55) and 0.58 (95% CI, 0.11 to 3.06), respectively (Fig C–H in [S1 Text](#)).

**Analysis of studies where all participants were ≥65 years.** Restricting the meta-analysis to only the 3 studies [[39–41](#)] that recruited participants with ≥65 years of age (total n = 58; analysed n = 50), effects were also broadly comparable to the primary analysis in the pattern of findings, with the exception of effect on all-cause AEs, which was no longer significant. The pooled IRRs for all-cause AEs (k = 3) were 2.13 (95% CI, 0.46 to 9.97) and treatment-related AEs (k = 2) was 2.80 (95% CI, 1.09 to 7.21), respectively. Pooled IRRs for all-cause (k = 3) and treatment-related SAEs (k = 2) were 1.80 (95% CI, 0.13 to 8.76) and 0.50 (95% CI, 0.03 to 7.99), respectively. Pooled RR for AE-related withdrawals (k = 3) and IRR for all deaths (k = 3) were 1.00 (95% CI, 0.23 to 4.41) and 0.42 (95% CI, 0.05 to 3.42), respectively.






**Fig 5. Forest plot of treatment-related serious adverse events: THC studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g005>

### THC:CBD combination studies

A total of 26 studies (5 crossover and 21 were parallel-arm; see Results in [S1 Text](#) and Table B in [S1 Text](#) for additional details) from 21 articles [[42,46,48,49,68–84](#)] reported on 1,965 patients [analysed 1,940; Total person-years of THC:CBD exposure: 394.29 person-years; Mean person-years of THC exposure (mean ± SD): 15.17 ± 28.20 person-years] on active and 1,887 (analysed 1,863) on placebo, ranging from 50 to 67 years in age (males: 0% to 80%). All studies used placebo as control. Two of the 26 included studies [[80](#)] investigated a combination of CBD and THCV, and we also examined the key effects after excluding these studies, which remained unchanged (please see Fig I–N in [S1 Text](#)). Results of meta-analysis for individual AEs after excluding these studies were identical to the results including all studies (as shown in Table D in [S1 Text](#)) and hence not shown.

Pooled IRRs for all-cause ( $k = 16$ ) and treatment-related ( $k = 9$ ) AEs from all RCTs were 1.58 (95% CI, 1.26 to 1.98) and 1.70 (95% CI, 1.24 to 2.33), respectively. Pooled IRRs for all-cause ( $k = 26$ ) and treatment-related ( $k = 21$ ) SAEs from all RCTs were 1.17 (95% CI, 0.99 to 1.39) and 1.19 (95% CI, 0.88 to 1.62), respectively. Pooled RR for AE-related withdrawals ( $k = 26$ ) and IRR for all deaths ( $k = 26$ ) from all RCTs were 1.40 (95% CI, 1.08 to 1.80) and 1.14 (95% CI, 0.89 to 1.46), respectively. Neither Egger test nor “Trim and fill” method indicated




**Fig 6. Forest plot of adverse event-related withdrawals: THC studies.** Numbers under the “Mean Age (yrs)” and “Withdrawals (n)” columns refer to the values in active and control intervention arms, respectively. The conditions listed are the disease conditions subgrouped into broader categories for meta-regression analyses purposes. They are: ND (dementia, Alzheimer disease, PD, Huntington disease, amyotrophic lateral sclerosis); MS; Cancer (cancer or chemotherapy-related anorexia, pain or nausea/vomiting, chemosensory alterations); and Other (type 2 diabetes mellitus, fibromyalgia, raised intraocular pressure, cervical dystonia, healthy, pancreatitis, obstructive sleep apnoea). MS, multiple sclerosis; NA, Not Available; ND, neurodegenerative disease; PD, Parkinson disease; RCT, randomised clinical trial; RR, risk ratio; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g006>

any significant effect of publication or other selection bias for any of the outcomes except for deaths (Fig O in S1 Text). For deaths as outcome, while Egger test was not significant, Trim and fill method indicated 9 missing studies, with the estimated effect becoming significant after their inclusion (1.33, 95% CI: 1.03 to 1.71;  $p = 0.027$ ,  $k = 35$ ). Where there was nonindependence of outcome data used in analyses, results of dependent meta-analyses were consistent with the results of independent meta-analyses.

**Effect of moderators.** Meta-regression analysis indicated that there was a significant effect of clinical condition on effect-size for treatment-related AEs [QM (df = 3) = 15.948,  $p = 0.01$ ] and AE-related withdrawals [QM (df = 3) = 8.987,  $p = 0.029$ ]. For treatment-related AEs, this was mainly related to a significantly higher effect-size in RCTs investigating pain conditions (regression coefficient = 0.393;  $p = 0.022$ ) and those investigating other conditions (regression coefficient = 1.263;  $p = 0.002$ ) compared to cancer conditions. For withdrawals, this was related to significantly higher effect-size in RCTs investigating pain conditions (regression coefficient = 0.816;  $p = 0.020$ ) and multiple sclerosis (regression coefficient = 0.675;  $p = 0.035$ )




**Fig 7. Forest plot of all deaths: THC studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g007>

compared to cancer conditions. Except these, moderators such as study design or type of intervention did not significantly influence estimated effects of THC:CBD combination treatment on any of the outcomes assessed.

**Effect of dose.** There was a significant effect of daily THC dose [QM (df = 1) = 4.554, p = 0.033] on AE-related withdrawals (Fig P in [S1 Text](#)) and a trend-level effect on all-cause




**Fig 8. Forest plot of all-cause adverse events: THC:CBD studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CBD, cannabidiol; CE, cannabis extract; IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g008>

[QM (df = 1) = 2.899, *p* = 0.089] and treatment-related AEs QM (df = 1) = 3.016, *p* = 0.082]. Exploratory analyses suggested that there was also a significant effect of CBD dose (QM (df = 1) = 4.539, *p* = 0.033) on all-cause AEs (Fig Q in [S1 Text](#)) and a trend-level effect [QM (df = 1) = 3.145, *p* = 0.076] on treatment-related AEs but no significant effect on withdrawals. Effects of dose of both THC and CBD were such that the higher their dose, the higher was the effect of THC:CBD combination treatment on withdrawals and AEs (all-cause and treatment-related). Except these, daily THC or CBD dose did not have any significant influence on the effects of THC:CBD combination treatment on SAEs or death.

**Common side effects.** Pooled IRRs of the most commonly reported AEs (Table D in [S1 Text](#)) suggested significantly higher incidence rate of nausea, vomiting, dry mouth, dizziness/light-headedness, somnolence/drowsiness, disorientation, fatigue, and visual symptoms in active compared to control arms.






**Fig 9. Forest plot of treatment-related adverse events: THC:CBD studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CBD, cannabidiol; CE, cannabis extract; IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g009>

**Analysis of studies where all participants were ≥50 years.** Restricting the meta-analysis to the 3 studies [42,68] that recruited participants with ≥50 years of age (total *n* = 60; analysed *n* = 52) was similar to the primary analysis in the pattern of findings, except for treatment-related AEs and AE-related withdrawals, which were no longer significant. The pooled IRRs for all-cause AEs (*k* = 3) and treatment-related AEs (*k* = 2) were 2.65 (95% CI, 1.49 to 4.71) and 4.73 (95% CI, 0.54 to 41.65), respectively. Pooled IRRs for all-cause (*k* = 3) and treatment-related SAEs (*k* = 3) were 1.00 (95% CI, 0.10 to 9.61) and 1.00 (95% CI, 0.10 to 9.61), respectively. Pooled RR for AE-related withdrawals (*k* = 3) and IRR for all deaths (*k* = 3) were 0.77 (95% CI, 0.18 to 3.22) and 1.00 (95% CI, 0.10 to 9.61), respectively (Fig R-W in S1 Text). None of these studies reported any SAEs (either treatment-related or all-cause) or death.

**Analysis of studies where all participants were ≥65 years.** As analysable data were available from only 1 study, this analysis was not carried out.






**Fig 10. Forest plot of all-cause serious adverse events: THC:CBD studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CBD, cannabidiol; CE, cannabis extract; IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g010>

### CBD studies


Four studies (3 crossover and 1 parallel-arm RCT) from 3 articles [50,80,85] reported on 43 patients (analysed 40) on active and 44 (analysed 41) on placebo, with age ranging from 53 to 59 years [53% to 100% males; person-years of total CBD exposure: 6.60 person-years; Mean person-years of CBD exposure (mean ± SD: 1.10 ± 1.23 person-years)]. Pooled IRR for all-cause AEs for all RCTs was 1.02 (95% CI, 0.90 to 1.16), based on data available only from the 2 crossover studies reporting on 3 different dosage conditions (Fig 14; Fig X in S1 Text). There was limited data to allow quantitative synthesis of other outcomes; however, there were no treatment-related SAEs, withdrawals, or death reported (see Results in S1 Text for qualitative synthesis). No analysable data was available from studies where all study participants were ≥50 years of age.



**Fig 11. Forest plot of treatment-related serious adverse events: THC:CBD studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CBD, cannabidiol; CE, cannabis extract; IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g011>

**Risk of bias.** A summary of the risk of bias of included studies are presented in Figs 15–19. Briefly, most RCTs reported adequate randomisation sequence generation and concealment, outcome objectiveness, and masking of outcome assessors; however, some studies had high risk of bias because of potential for unmasking of participants and study personnel and selective reporting of the safety outcome. Most studies reported objective outcome assessments; however, only 60% of studies reported that outcome assessors had been appropriately blinded. A total of 45% of studies did selective reporting, i.e., they did not report data for all the safety outcomes (AEs and SAEs) in the trial and reported them when they occurred more than once or were more common or when they had occurrence above a certain threshold (1% to 10%). The authors judged 33 (55%) trials at low risk of bias, 20 (33%) trials at unclear risk of bias, and 7 (12%) trials to have high risk of bias for safety outcome reporting (Fig 15–19). Overall, 36 trials were judged to be of moderate quality, of which 15 (42%) trials reported all




**Fig 12. Forest plot of all withdrawals: THC:CBD studies.** Numbers under the “Mean Age (yrs)” and “Withdrawals (n)” columns refer to the values in active and control intervention arms, respectively. The conditions listed are the disease conditions subgrouped for meta-regression analyses purposes are: MS; MND; pain (neuropathic pain, rheumatoid arthritis), cancer (cancer or chemotherapy-related anorexia, pain or nausea/vomiting), diabetes mellitus, COPD, HC, levodopa-induced dyskinesia in PD. CBD, cannabidiol; COPD, chronic obstructive pulmonary disease; HC, healthy controls; MS, multiple sclerosis; MND, motor neurone disease; PD, Parkinson disease; RCT, randomised clinical trial; RR, risk ratio; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g012>

AEs and SAEs. Ten (56%) of moderate-quality trials of THC only intervention reported all AEs and SAEs.

### Discussion

In this systematic review and meta-analysis, we investigated the safety and tolerability of medicinal cannabinoids in older adults by pooling data from double-blind RCTs with a reported mean participant age of 50 years and over. We hypothesized that compared to control treatments, all 3 categories of CBMs will be associated with a greater incidence of AEs, but no greater incidence of SAEs, death, or risk of withdrawal from study. We also expected a direct




**Fig 13. Forest plot of all deaths: THC:CBD studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CBD, cannabidiol; CE, cannabis extract; IRR, incident rate ratio; RCT, randomised clinical trial; THC, delta-9-tetrahydrocannabinol.

<https://doi.org/10.1371/journal.pmed.1003524.g013>

relationship between the dose of THC used in THC-containing CBMs and the risk of adverse outcomes. We found that generally moderate to high-quality evidence (about 60% studies) suggests that as hypothesized, treatment with THC-containing medications (THC alone and THC:CBD combination) was associated on average with significantly higher incidence of all-cause and treatment-related AEs compared to control treatments. Further, consistent with our hypotheses, the average incidence rates of serious AEs (all-cause and treatment-related) and death were not significantly greater under CBMs compared to controls in studies using THC with or without CBD. However, contrary to expectation, significantly higher risk of withdrawal related to AEs was noted on average for studies using THC:CBD combination, though this was not observed in studies that used THC without CBD. In contrast, generally low-quality evidence (about 67% of studies) suggests that CBD alone may not significantly increase the incidence rate of all cause of AEs. Qualitative synthesis of data on treatment-related AEs, SAEs, deaths, and withdrawals from study also did not suggest any increase associated with CBMs containing CBD alone. In terms of relationship between THC dose and adverse





**Fig 14. Forest plot of all-cause adverse events: CBD studies.** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. Overall study quality GRADE [31] is reported in Tables 1–3 and Results in S1 Text. Risk of bias estimates are reported in Figs 15–19. CBD, cannabidiol; GRADE, Grading of Recommendations Assessment, Development, and Evaluation; IRR, incident rate ratio; RCT, randomised clinical trial.


<https://doi.org/10.1371/journal.pmed.1003524.g014>

outcomes, as hypothesized, we found a direct relationship between daily THC dose and all-cause AEs and AE-related withdrawals in THC studies and between THC dose and AE-related withdrawals in THC:CBD studies. Contrary to our expectations, we did not find any association between THC dose and the other outcomes investigated, such as SAEs and deaths. In addition, exploratory analysis showed that CBD dose also had a significant direct relationship with all-case AEs and a strong trend-level association with treatment-related AEs in RCTs using THC-CBD combination.


Additional analyses restricted to only those studies in which all participants were ≥50 years of age or ≥65 years of age, where this was feasible, indicated a pattern of findings broadly comparable with the results of our main analyses including all studies. However, the effect of CBMs containing THC but no CBD on all-cause AEs and CBMs containing THC:CBD on treatment-related AEs and AE-related withdrawals were no longer significant, which likely reflects the lower power of these analyses which included a maximum of 4 studies in any analysis.

Collectively, our results from studies that included participants with a mean age ≥ 50 years may suggest that older adults are at significantly greater risk of both treatment-related and all-cause AEs from CBMs containing THC, but not using CBMs without THC. Despite the greater risk of AEs, older adults receiving CBMs do not seem to be at a significantly greater risk of





**Fig 15. Risk of bias (THC studies).** Authors' judgements about each risk of bias domain for each individual study.  
<https://doi.org/10.1371/journal.pmed.1003524.g015>



**Fig 16. Summary of risk of bias (THC studies).** Authors' judgements about each risk of bias domain reported as percentages across included studies.

<https://doi.org/10.1371/journal.pmed.1003524.g016>

more serious consequences such as SAEs or death. Greater risk of withdrawal from study in those receiving CBMs containing THC:CBD combination but not in those containing THC or CBD without the other cannabinoid is an unexpected finding. While we did not compare these 2 effects in a systematic manner, if true, this may suggest that combination of these cannabinoids may be less acceptable in this age group, who already may be receiving multiple treatments for comorbidities. What may underlie these effects is much less clear. One may speculate that this may be a result of the generally higher THC dose employed in THC:CBD studies compared to the THC studies, which may have made participants in the former group of studies more susceptible to AEs and withdrawal as a result compared to those receiving THC. This is supported by approximately double the median dose of THC used in THC:CBD studies [median and interquartile range: 10.3(10.2 to 21.6) mg/day in crossover and 20.8(14.6 to 25) mg/day in parallel-arm RCTs] as opposed to THC studies [median and interquartile range: 5(2.25 to 9 mg/day in crossover and 10(4.8 to 25) mg/day in parallel-arm RCTs]. In





Fig 17. Risk of bias (THC:CBD studies). Authors' judgements about each risk of bias domain for each individual study.

<https://doi.org/10.1371/journal.pmed.1003524.g017>



**Fig 18. Summary of risk of bias (THC:CBD studies).** Authors' judgements about each risk of bias domain reported as percentages across included studies.

<https://doi.org/10.1371/journal.pmed.1003524.g018>

accordance with this hypothesis, there was a nominally higher incidence of AEs in general in those studies using both cannabinoids in combination compared to THC studies. However, one cannot completely rule out the possibility that greater risk of withdrawal from study in those receiving CBMs containing THC:CBD combination but not in those containing THC or CBD without the other cannabinoid, as observed here, is purely by chance. Further, THC dose was significantly associated with withdrawal in both study groups using THC-containing medications. In contrast, CBD dose was only associated with AEs but not withdrawal in those using THC:CBD combination. However, whether these results suggest that lower acceptability of THC:CBD combination is a tolerability issue in older patients because of combination of cannabinoids rather than being related to the dose of CBD remains to be formally tested. It is also worth noting that the evidence base of studies that used CBMs without THC in older adults and found it to be well tolerated is relatively sparse. Whether this may underlie the absence of significantly greater risk of all-cause AEs as a result of treatment with CBMs

		Risk of bias domains								
		sequence generation	allocation concealment	blinding personnel	blinding outcome assessment	selective outcome reporting	attrition	selective reporting	Other biases	Stopped early
Study	Consroe et al. 1991	?	?	+	+	+	+	+	+	+
	Jadoon et al. 2016	+	+	+	?	+	+	x	+	+
	Tomida et al. 2006	?	?	+	?	?	+	+	+	+

**Fig 19. Risk of bias (CBD studies).** Authors' judgements about each risk of bias domain for each individual study.

<https://doi.org/10.1371/journal.pmed.1003524.g019>

without THC compared to controls, contrary to our expectation, remains to be tested. However, more generally, lack of significant AEs associated with CBD treatment alone in the present study is consistent with other analyses suggesting a good tolerability profile of CBD alone in younger age groups [86,87]. Nevertheless, collectively, these results suggest that although the incidence of side effects is greater with THC-containing products, CBMs investigated in these studies are generally safe. Their overall tolerability profile also seems to be generally acceptable to patients, though combination of THC and CBD may be less so. It is important to note that although the general pattern of results of analyses restricted to studies where all participants were  $\geq 50$  years of age or  $\geq 65$  years of age were broadly comparable with results from our main analyses of studies with mean participant age  $\geq 50$  years, they were not identical, indicating that effects of CBMs may be different when focused exclusively on adults over 65 or 75 years of age.

Although we did observe a significant association between THC dose and all-cause AEs in THC studies, this was not present in THC:CBD studies. Whether this reflects independent evidence that CBD may mitigate some of the adverse effects of THC, such as on cognition, behaviour, autonomic, and cardiovascular function [15,25–28] and therefore may have obscured any dose–response relationship with AEs in THC:CBD studies, remains unclear. The present study was not designed to formally test this hypothesis, which needs independent examination.

Finally, we identified a number of specific AEs associated with CBMs containing THC: CBD combination or THC without CBD. Of the specific AEs, significantly increased incidence of dizziness/light-headedness, drowsiness, disorientation, and impaired mobility/balance/coordination is worth noting, in light of higher risk of falls in this age group, which may be further exacerbated by these treatments [88,89].

Pooled estimates of all-cause AEs under both THC and THC:CBD and of treatment-related AEs and AE-related withdrawals under THC:CBD as reported herein need to be considered

carefully in light of high levels of heterogeneity (discussed further below) in the studies as evident from the estimated  $I^2$  statistic ( $>32\%$  in all these estimates). Therefore, we have also reported the prediction intervals along with the 95% confidence intervals in the forest plots for the key pooled estimates reported herein. These prediction intervals indicate that while the average effects of CBM treatments on these outcomes are significant (as evident from the 95% confidence intervals), the range of predicted effects across different study settings that may be observed in a new study may span a wider range of effect sizes than indicated by the 95% confidence intervals. Some of these predicted effect-sizes in new studies are likely to be  $<1$ , indicating that the risk of these outcomes with THC or THC:CBD treatment may not be higher than under control treatment condition.

Previous reviews of adverse consequences of treatment with CBMs have either been qualitative [12,19,20], did not specifically focus on older adults [16,17], or did not consider the effects of THC, CBD, or their combination separately [12,16,17]. Even in studies that have pooled the adverse event data quantitatively, this has been done based on specific formulation rather than on the basis of their cannabinoid content [16,17]. Further, they have generally reported summary effect-size estimates (such as odds ratio) that do not take into account person-years of treatment, an important consideration in quantitative synthesis of outcomes from RCTs [16,17]. As in previous reviews, we found that not all studies have published all the AEs and SAEs and that there is lack of evidence of the safety, tolerability, and acceptability specifically in older patients. Nevertheless, our findings are consistent with previous reviews which found dizziness to be the most common nonserious AE with CBM treatment [12,16,17]. These results are also consistent with a prospective observational study of 901 people above 65 years of age ( $74.5 \pm 7.5$  years), who received medical cannabis from January 2015 to October 2017 in a specialized medical cannabis clinic, 31.7% reported at least 1 AE due to the treatment after 6 months, with dizziness (9.7%) and dry mouth (7.1%) as the commonest AEs [21]. Another similar study of 184 people ( $81.2 \pm 7.5$  years of age) from April 2017 to October 2018 showed 33.6% with AEs and dizziness (12.1%) and sleepiness and fatigue (11.2%) as the commonest AEs [90]. A previous systematic review of the efficacy and safety of medicinal cannabinoids which focused only on older people considered 5 controlled trials with THC ( $n = 3$ ) and oral THC:CBD ( $n = 2$ ) [12]. However, this review did not provide summary estimates (effect sizes) due to high heterogeneity among the included studies, lack of available data on means and standard deviations per treatment group, and very small sample sizes [12]. Therefore, only qualitative and descriptive summaries were provided. Studies included in this review reported dizziness, euphoria, drowsiness, confusion, and disorientation as the common adverse effects [12], consistent with our report. An earlier systematic review of AEs of medicinal cannabinoids for all ages by Wang and colleagues [17] included 23 RCTs published between 1966 and late 2007 and analysed the effect of oral THC and THC:CBD oral and spray formulations. However, they did not include data from studies examining the synthetic cannabinoid nabilone and found no evidence of higher incidence of SAEs after a median of 2 weeks use compared with a control group, regardless of the age of individuals (RR 1.04, 95% CI 0.78 to 1.30). Wang and colleagues [17] reported that respiratory, gastrointestinal, and nervous system disorders were the most frequently reported categories of SAEs, and dizziness was the most commonly reported nonserious AE (15.5%) among people exposed to cannabinoids. However, unlike the present review wherein we have used GRADE criteria to assess methodological quality of included RCTs, Wang and colleagues used a scale reported in a manuscript by Jadad and colleagues [91] to assess methodological quality of RCTs. As the scale from Jadad and colleagues does not adequately assess the quality of safety reporting in RCTs, most studies were rated as of good quality in the review by Wang and colleagues, despite their poor reporting of safety [17].



However, our results are not consistent with a more recent systematic review and meta-analysis by Whiting and colleagues [16], which included studies investigating all age groups and showed that SAEs and AE-related withdrawal are generally higher in those treated with CBMs. In contrast, we have shown that SAEs are not increased in older adults consistent with other reviews [12,17]. Whiting and colleagues [16] found pooled effect-size for any AE (OR 3.03, 95% CI 2.42 to 3.80), SAE (OR 1.41, 95% CI 1.04 to 1.92), and withdrawal due to AE (OR 2.94, 95% CI 2.18 to 3.96), and Wang and colleagues [17] found significantly higher nonserious AEs (RR 1.86, 95% CI 1.57 to 2.21) in those who received CBMs compared to control groups. In contrast to Whiting and colleagues, we have shown that increase in withdrawal may be more nuanced and more in those receiving THC:CBD combination and not for other types of CBMs.

It is difficult to compare our pooled effect-size with previous reviews due to a number of reasons. Firstly, Whiting reported odds ratios and Wang and colleagues reported relative risk, wherein we have reported IRR. In the present context, the risk of AEs or SAEs in either study arm are unlikely to be constant over time. Therefore, IRR, which takes into consideration person-years of treatment and is a ratio of the incidence rate in the experimental treatment group to that in the control treatment group, is more meaningful and appropriate in contrast to the odds ratio, which is a ratio of the odds of an event in the experimental group to that in the control group and are not easily interpretable. Secondly, we carried out a pooled analysis for individual interventions unlike the other reviews, with Wang and colleagues also excluding studies using Nabilone [17]. Finally, the 2 previous reviews that reported summary estimates included all age groups and were not specific for older people unlike here.

Our results extend previous literature by showing that although at significantly higher risk of adverse events from CBMs containing THC, but not from those without THC, CBMs are generally safe in older adults, who typically experience comorbid health conditions [92] and receive polypharmacy [93]. Also, CBMs are generally acceptable as long as they are not a combination of THC and CBD.

Results presented here may also need to be considered against the side effect profile of common treatments for the clinical conditions investigated in studies included in the meta-analyses reported here, as safety and tolerability profile of a potential new treatment relative to existing alternatives is an important consideration in the context of prescribing. Studies included patients with Alzheimer disease, Parkinson disease, Huntington disease, amyotrophic lateral sclerosis, multiple sclerosis, motor neurone disease, neuropathic pain, cancer (cancer or chemotherapy-related anorexia, pain or nausea/vomiting), type 2 diabetes mellitus, chronic obstructive pulmonary disease, fibromyalgia, raised intraocular pressure, cervical dystonia, pancreatitis, and obstructive sleep apnoea. As would be expected, these conditions are typically treated with a wide range of pharmacological treatments (opioid analgesics, nonsteroidal anti-inflammatory, antiepileptics, benzodiazepines, psychotropics, cholinesterase inhibitors, glutamate antagonists, dopamine agonists, immunosuppressants, muscle relaxants, etc.) with a varied side effect profile. While a systematic comparison of side effect profile of CBMs and these other treatments would be useful and can helpfully inform prescribing decisions, this was outside the scope of the present review and was not examined here. Therefore, future studies may need to investigate this in a systematic manner. As we pooled data from randomised controlled comparisons, as opposed to observational studies without a control arm, they are very unlikely to have confounded the results presented herein.

## Strengths and limitations

Our review and meta-analysis are limited by a number of methodological weaknesses, some of which stem from the design and analytic approach of the present study and others which are

related to weaknesses in the studies that were included in the present analysis. One of the key limitations inherent in the meta-analytic approach that also applies to the present study relates to the issue of heterogeneity in the pooled data [94]. Studies pooled in our analyses focused on patients with different clinical conditions and indications, used different doses, formulations, and routes of administration of the study drug for different treatment periods and employed different study designs (crossover versus parallel-arm RCTs). Further, although we attempted to control for patients from widely disparate age groups taking part in the trial by setting an inclusion criterion of mean age of 50 years and over as the cut-off, studies included in our main analyses still involved data from people below age 50 and studies also varied in terms of sex distribution. This is reflected in the heterogeneity estimates reported in our results. In order to address this wide-ranging heterogeneity and yet pool the data in a meaningful way, we employed a random-effects model in our analyses, which assumes that there will be variability in the observed estimates of treatment effects across studies, both as a result of real differences in the effect of treatment between studies as well as by chance because of sampling variability. As a result, pooled estimates reported are not precise (as may be evident from the wide confidence intervals) and should not be considered as such. Effects reported here represent an average effect of CBM treatments on safety and tolerability outcomes investigated rather than an effect that is common across studies. As the effects may be different within an individual study, we also report a prediction interval for our key reported outcomes, in order to give an estimate of the range of predicted effects across study settings that may be observed in a new study. Further, we also carried out subgroup (e.g., crossover and parallel-arm RCTs separately) and meta-regression analyses to examine the sources of heterogeneity for THC and THC:CBD studies. These analyses did not suggest that study design or type of intervention had any significant effect on the outcomes assessed, though they suggest that effects vary, sometimes significantly, across clinical conditions investigated in the RCTs. They were more pronounced in AE-related withdrawals and treatment-related AEs in THC:CBD studies and less so for all-cause AEs in studies investigating THC as a treatment. Dose of study drug also seemed to underlie some of the heterogeneity observed in all-cause and treatment-related AEs and AE-related withdrawals in the THC and THC:CBD studies. Another limitation relates to the fact that we were not able to systematically examine the sources of heterogeneity for CBD alone studies as there were fewer studies than recommended for such analyses. One other limitation of the present meta-analysis relates to our focus on studies reporting on participants aged 50 years and over.

Further, we included studies in which the mean age of study participants was  $\geq 50$  years (although the studies also included many participants who were  $< 50$  years of age). As the cut-off employed by us differs from the conventional threshold of 65 years for “elderly” [95], this may be considered as a limitation. However, this was chosen as the clinical conditions (diabetes, cancer, neurodegenerative disorders, etc.) for which CBMs are often considered become more common from around this age. This is also a period characterised by multimorbidities, polypharmacy, and age-related bodily changes that may affect pharmacokinetics [95] and tolerability of medications. Further, in order to address the limitation that many participants in the included studies were  $< 50$  years of age, we also carried out sensitivity analysis that included studies where all participants were  $\geq 50$  years of age. The results of these analyses suggest that the pooled effect-sizes were generally in the same direction as that reported from the larger set of studies, though the confidence intervals were wider, as may be expected. We have also reported the number of studies which have actually studied individuals with age  $\geq 65$  years or  $\geq 75$  years. As evident from this, there is a very limited set of studies that have exclusively focused on people at these ages. Therefore, the present meta-analysis highlights the need for studies that may need to focus on people over 65. However, given the age range as well as

median and interquartile range of the mean ages of study participants included in the studies that constitute our meta-analysis, it is clear that people over 65 and 75 years are currently being recruited into studies of CBMs for various indications. As individual RCTs are often not powerful enough to unravel patterns of side effects and given the growing use of CBMs in the elderly and general perception of them being safe, it is particularly important to summarize currently available evidence to help inform about the safety and tolerability profile of CBMs in those aged 50 years and over rather than wait for the evidence base focusing only on  $\geq 65$  years to mature. In the fullness of time, future attempts at evidence synthesis need to focus only on studies of people  $\geq 65$  years when a sufficient number of studies have accumulated.

The other main source of limitation stems from methodological limitations in the included trials as identified during quality assessment [30,31], specifically pertaining to selective outcome reporting, and inadequate description of methods of randomization, allocation concealment, and blinding. Additionally, many included RCTs investigated only modestly sized samples [41,42,44,50,57]. Small samples render studies particularly underpowered when estimating serious and less serious adverse outcomes, which by their nature may not occur frequently. It is in this context that the present report addresses an important gap in extant evidence by systematic quantitative synthesis of RCT data following existing recommendations [30,31] to provide estimates from a larger pool of patients. Further, our analyses suggest that publication or other selection biases are unlikely to have influenced the pooled estimates reported here.

Unlike in previous meta-analyses, which reported summary effects separately based on indications, we pooled safety and tolerability data in older adults across a broad range of indications. While this may have added to the heterogeneity of the data synthesized, it allowed us to comprehensively estimate separately the effects of 3 broad categories of cannabinoid-based interventions, i.e., THC only, THC:CBD combination, and CBD only, something that has not been done before. This is a key strength of the present approach, given the reported opposite effects of different cannabinoids [4,15] that argue against data being combined. Another important strength of the present report relates to the analysis of the effects of moderators to examine the extent to which they may have influenced results, in particular relationship with cannabinoid doses used.

Studies evaluated various routes of CBM administration (oral capsules, tablets, sublingual spray, oromucosal spray). Also, not all studies compared CBMs with placebo, with 4 studies using active control treatments [43,45,54,62]. While all of these may have resulted in a very heterogeneous set of included studies, we used a random-effects model to mitigate these effects. Further, heterogeneity did not seem to significantly affect any of our estimates other than all-cause AEs, giving further confidence in the results reported. Nevertheless, we have also reported prediction intervals in addition for our key reported outcomes to give an estimate of the range of predicted effects across different study settings. Finally, another important potential limitation of the present study relates to the fact that we did not investigate the efficacy of CBMs in older adults. As outcome measures used to index efficacy vary widely between clinical conditions and there is a relative paucity of studies investigating a particular clinical condition, there are not enough data for any quantitative synthesis of efficacy of different CBMs in older adults to be meaningful just yet.

Clinical efficacy is one of the foremost considerations in addition to patient choice and safety/tolerability of interventions when prescribing in clinical practice. While the present study summarizes current evidence regarding safety/tolerability of CBMS, there is limited efficacy evidence for most clinical indications for which CBMS have been used in older people. Therefore, there is a pressing need for efficacy studies in specific indications where there is proof of concept or rationale for use of CBMs in older people. With regard to CBMs, potential

for drug–drug interaction in light of effect on cytochrome p450 enzymes is a major concern in the context of treating older patients [13]. However, few studies have examined this, an important likely determinant of tolerability and dose adjustment, and therefore worthy of investigation in future studies.

Complete reporting of safety/tolerability data as well as improved trial designs incorporating robust methods for allocation concealment, masking of participant and outcome assessors are further important considerations for future trials. Using well-powered samples, such studies need to focus on safety, tolerability, as well as efficacy of different categories of CBMs, in particular CBD on its own, a relatively less investigated CBM in older people.

## Conclusions

Results of the present study using data from RCTs with mean participant age  $\geq 50$  years suggest that although THC-containing CBMs are associated with side effects in those aged 50 years and over, in general, CBMs are safe and acceptable treatments in older adults, with a caveat that THC:CBD combinations may be less so at least in dose ranges used in studies thus far. However, tolerability may be different in adults over 65 or 75 years of age, and robust evidence of efficacy of different CBMs for specific indications is needed before they may be used in routine practice in older adults.

## Supporting information

### S1 PRISMA Checklist. PRISMA checklist.

(DOC)

**S1 Text. Supporting Methods, Results, Tables and Figures.** Table A in [S1 Text](#). Summary of randomised controlled trials of THC in older adults for studies with participants with mean age  $\geq 50$  years ( $N = 30$ ) and studies with participants with age  $\geq 50$  years ( $N = 4$ , 13%; one  $\geq 50$  years; two  $\geq 65$  years; one  $\geq 75$  years). Table B in [S1 Text](#). Summary of randomised controlled trials of THC:CBD combination in older adults for studies with participants with mean age  $\geq 50$  years ( $N = 26$ ) and studies with participants recruited with age  $\geq 50$  years ( $N = 3$ , 11.5%; two  $\geq 50$  years; one  $\geq 65$  years). Table C in [S1 Text](#). Summary estimates (incident rate ratio, IRR) from meta-analysis for the most commonly reported adverse events (AEs): IRR of participants experiencing AE with cannabinoid (THC) compared to placebo or active control condition. Table D in [S1 Text](#). Summary estimates (incident rate ratio, IRR) from meta-analysis for the most commonly reported adverse events (AEs): IRR of participants experiencing AE with cannabinoid (THC:CBD combination) compared to placebo or active control condition. Table E in [S1 Text](#). Characteristics of unpublished randomised trials of cannabinoids in older adults. **Fig A in S1 Text. Funnel plots for all tolerability and safety outcomes: THC studies.** (a) All-cause adverse events (AEs); (b) Treatment-related AEs; (c) All-cause serious adverse events (AAEs); (d) Treatment-related SAEs; (e) AE-related withdrawals; (f) deaths. **Fig B in S1 Text.** THC dose-related withdrawals in THC studies. **Fig C in S1 Text.** Forest plot of all-cause adverse events: THC studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. **Fig D in S1 Text.** Forest plot of treatment-related serious adverse events: THC studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. **Fig E in S1 Text.** Forest plot of all-cause serious adverse events: THC studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. **Fig F in S1 Text.** Forest plot of treatment-



related serious adverse events: THC studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. Fig G in [S1 Text](#). Forest plot of adverse event-related withdrawals: THC studies (participants with  $\geq 50$  years of age). Numbers under the “Mean Age (yrs)” and “Withdrawals (n)” columns refer to the values in active and control intervention arms, respectively. The conditions listed are the disease conditions subgrouped into broader categories for meta-regression analyses purposes. They are: neurodegenerative disorders (ND) (dementia, Alzheimer disease, Parkinson disease (PD), Huntington disease, amyotrophic lateral sclerosis); multiple sclerosis (MS); Cancer (cancer or chemotherapy-related anorexia, pain or nausea/vomiting, chemosensory alterations); and Other (type 2 diabetes mellitus, fibromyalgia, raised intraocular pressure, cervical dystonia, healthy, pancreatitis, obstructive sleep apnoea). **Fig H in [S1 Text](#). Forest plot of all deaths: THC studies (participants with  $\geq 50$  years of age).** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. **Fig I in [S1 Text](#). Forest plot of all-cause adverse events: THC:CBD studies (excluding THCv).** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig J in [S1 Text](#). Forest plot of treatment-related adverse events: THC:CBD studies (excluding THCv). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig K in [S1 Text](#). Forest plot of all-cause serious adverse events: THC:CBD studies. (excluding THCv). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig L in [S1 Text](#). Forest plot of treatment-related serious adverse events: THC:CBD studies (excluding THCv). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. **Fig M in [S1 Text](#). Forest plot of all withdrawals: THC:CBD studies (excluding THCv).** Numbers under the “Mean Age (yrs)” and “Withdrawals (n)” columns refer to the values in active and control intervention arms, respectively. The conditions listed are the disease conditions subgrouped for meta-regression analyses purposes are: multiple sclerosis (MS); motor neurone disease (MND); pain (neuropathic pain, rheumatoid arthritis), cancer (cancer or chemotherapy-related anorexia, pain or nausea/vomiting), diabetes mellitus, chronic obstructive pulmonary disease (COPD), healthy controls (HC), levodopa-induced dyskinesia in Parkinson disease (PD). **Fig N in [S1 Text](#). Forest plot of all deaths: THC:CBD studies (excluding THCv).** Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. **Fig O in [S1 Text](#). Funnel plots for all tolerability and safety outcomes: THC:CBD studies.** (a) All-cause adverse events (AEs); (b) Treatment-related AEs; (c) All-cause serious adverse events (AAEs); (d) Treatment-related SAEs; (e) AE-related withdrawals; (f) deaths. Fig P in [S1 Text](#). THC dose-related withdrawals in THC:CBD studies. Fig Q in [S1 Text](#). CBD dose-related all-cause AEs in THC:CBD studies. Fig R in [S1 Text](#). Forest plot of all-cause adverse events: THC:CBD studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig S in [S1 Text](#). Forest plot of treatment-related adverse events: THC:CBD studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig T in [S1 Text](#). Forest plot of all-cause serious adverse events: THC:CBD studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig U in [S1 Text](#). Forest plot of



treatment-related serious adverse events: THC:CBD studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig V in [S1 Text](#). Forest plot of all withdrawals: THC:CBD studies (participants with  $\geq 50$  years of age). Numbers under the “Mean Age (yrs)” and “Withdrawals (n)” columns refer to the values in active and control intervention arms, respectively. The conditions listed are the disease conditions subgrouped for meta-regression analyses purposes are: multiple sclerosis (MS); motor neurone disease (MND); pain (neuropathic pain, rheumatoid arthritis), cancer (cancer or chemotherapy-related anorexia, pain or nausea/vomiting), diabetes mellitus, chronic obstructive pulmonary disease (COPD), healthy controls (HC), levodopa-induced dyskinesia in Parkinson disease (PD). Fig W in [S1 Text](#). Forest plot of all deaths: THC:CBD studies (participants with  $\geq 50$  years of age). Numbers under the “Subjects (n)” column refer to analysed participants from the active and control intervention arms, respectively. CE refers to cannabis extract. Fig X in [S1 Text](#). Funnel plot for all-cause adverse events (AEs): CBD studies. (DOCX)

## Acknowledgments

None.

The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, PUK or the Department of Health and Social Care.

## Author Contributions

**Conceptualization:** Latha Velayudhan, Sagnik Bhattacharyya.

**Data curation:** Latha Velayudhan, Katie McGoohan.

**Formal analysis:** Sagnik Bhattacharyya.

**Methodology:** Latha Velayudhan, Katie McGoohan, Sagnik Bhattacharyya.

**Software:** Sagnik Bhattacharyya.

**Supervision:** Sagnik Bhattacharyya.

**Validation:** Latha Velayudhan.

**Writing – original draft:** Latha Velayudhan.

**Writing – review & editing:** Katie McGoohan, Sagnik Bhattacharyya.

## References

1. Hazekamp A, Ware MA, Muller-Vahl KR, Abrams D, Grotenhermen F. The medicinal use of cannabis and cannabinoids—an international cross-sectional survey on administration forms. *J Psychoactive Drugs*. 2013; 45(3):199–210. Epub 2013/11/02. <https://doi.org/10.1080/02791072.2013.805976> PMID: [24175484](https://pubmed.ncbi.nlm.nih.gov/24175484/).
2. Kaskie B, Ayyagari P, Milavetz G, Shane D, Arora K. The Increasing Use of Cannabis Among Older Americans: A Public Health Crisis or Viable Policy Alternative? *Gerontologist*. 2017; 57(6):1166–72. <https://doi.org/10.1093/geront/gnw166> PMID: [28077451](https://pubmed.ncbi.nlm.nih.gov/28077451/). Language: English. Entry Date: 20171129. Revision Date: 20181203. Publication Type: Article.
3. Hazekamp A, Heerdink ER. The prevalence and incidence of medicinal cannabis on prescription in The Netherlands. *Eur J Clin Pharmacol*. 2013; 69(8):1575–80. Epub 2013/04/17. <https://doi.org/10.1007/s00228-013-1503-y> PMID: [23588562](https://pubmed.ncbi.nlm.nih.gov/23588562/).
4. Pertwee RG. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol*. 2008; 153

- (2):199–215. Epub 2007/09/11. <https://doi.org/10.1038/sj.bjp.0707442> PMID: [17828291](https://pubmed.ncbi.nlm.nih.gov/17828291/); PubMed Central PMCID: PMC2219532.
5. Bhattacharyya S, Fusar-Poli P, Borgwardt S, Martin-Santos R, Nosarti C, O'Carroll C, et al. Modulation of mediotemporal and ventrostriatal function in humans by Delta9-tetrahydrocannabinol: a neural basis for the effects of Cannabis sativa on learning and psychosis. *Arch Gen Psychiatry*. 2009; 66(4):442–51. Epub 2009/04/08. <https://doi.org/10.1001/archgenpsychiatry.2009.17> PMID: [19349314](https://pubmed.ncbi.nlm.nih.gov/19349314/).
  6. Babalonis S, Haney M, Malcolm RJ, Lofwall MR, Votaw VR, Sparenborg S, et al. Oral cannabidiol does not produce a signal for abuse liability in frequent marijuana smokers. *Drug Alcohol Depend*. 2017; 172:9–13. Epub 2017/01/15. <https://doi.org/10.1016/j.drugalcdep.2016.11.030> PMID: [28088032](https://pubmed.ncbi.nlm.nih.gov/28088032/); PubMed Central PMCID: PMC5361620.
  7. Burstein S. Cannabidiol (CBD) and its analogs: a review of their effects on inflammation. *Bioorg Med Chem*. 2015; 23(7):1377–85. Epub 2015/02/24. <https://doi.org/10.1016/j.bmc.2015.01.059> PMID: [25703248](https://pubmed.ncbi.nlm.nih.gov/25703248/).
  8. Bhattacharyya S, Wilson R, Appiah-Kusi E, O'Neill A, Brammer M, Perez J, et al. Effect of Cannabidiol on Medial Temporal, Midbrain, and Striatal Dysfunction in People at Clinical High Risk of Psychosis: A Randomized Clinical Trial. *JAMA Psychiatry*. 2018; 75(11):1107–17. Epub 2018/09/01. <https://doi.org/10.1001/jamapsychiatry.2018.2309> PMID: [30167644](https://pubmed.ncbi.nlm.nih.gov/30167644/); PubMed Central PMCID: PMC6248101
  9. Leweke FM, Mueller JK, Lange B, Rohleder C. Therapeutic Potential of Cannabinoids in Psychosis. *Biol Psychiatry*. 2016; 79(7):604–12. Epub 2016/02/08. <https://doi.org/10.1016/j.biopsych.2015.11.018> PMID: [26852073](https://pubmed.ncbi.nlm.nih.gov/26852073/).
  10. Allan GM, Finley CR, Ton J, Perry D, Ramji J, Crawford K, et al. Systematic review of systematic reviews for medical cannabinoids: Pain, nausea and vomiting, spasticity, and harms. *Can Fam Physician*. 2018; 64(2):e78–e94. Epub 2018/02/17. PMID: [29449262](https://pubmed.ncbi.nlm.nih.gov/29449262/); PubMed Central PMCID: PMC5964405.
  11. Fraguas-Sánchez AI, Torres-Suárez AI. Medical Use of Cannabinoids. *Drugs*. 2018; 78(16):1665–703. Epub 2018/10/31. <https://doi.org/10.1007/s40265-018-0996-1> PMID: [30374797](https://pubmed.ncbi.nlm.nih.gov/30374797/).
  12. van den Elsen GA, Ahmed AI, Lammers M, Kramers C, Verkes RJ, van der Marck MA, et al. Efficacy and safety of medical cannabinoids in older subjects: a systematic review. *Ageing Res Rev*. 2014; 14:56–64. Epub 2014/02/11. <https://doi.org/10.1016/j.arr.2014.01.007> PMID: [24509411](https://pubmed.ncbi.nlm.nih.gov/24509411/).
  13. Mahvan TD, Hilaire ML, Mann A, Brown A, Linn B, Gardner T, et al. Marijuana Use in the Elderly: Implications and Considerations. *Consult Pharm*. 2017; 32(6):341–51. Epub 2017/06/10. <https://doi.org/10.4140/TCP.n.2017.341> PMID: [28595684](https://pubmed.ncbi.nlm.nih.gov/28595684/).
  14. Hulshof TA, Zuidema SU, Ostelo RW, Luijendijk HJ. The Mortality Risk of Conventional Antipsychotics in Elderly Patients: A Systematic Review and Meta-analysis of Randomized Placebo-Controlled Trials. *J Am Med Dir Assoc*. 2015; 16(10):817–24. Epub 2015/05/03. <https://doi.org/10.1016/j.jamda.2015.03.015> PMID: [25933724](https://pubmed.ncbi.nlm.nih.gov/25933724/).
  15. Bhattacharyya S, Morrison PD, Fusar-Poli P, Martin-Santos R, Borgwardt S, Winton-Brown T, et al. Opposite effects of delta-9-tetrahydrocannabinol and cannabidiol on human brain function and psychopathology. *Neuropsychopharmacology*. 2010; 35(3):764–74. Epub 2009/11/20. <https://doi.org/10.1038/npp.2009.184> PMID: [19924114](https://pubmed.ncbi.nlm.nih.gov/19924114/); PubMed Central PMCID: PMC3055598.
  16. Whiting PF, Wolff RF, Deshpande S, Di Nisio M, Duffy S, Hernandez AV, et al. Cannabinoids for Medical Use: A Systematic Review and Meta-analysis. *Jama*. 2015; 313(24):2456–73. Epub 2015/06/24. <https://doi.org/10.1001/jama.2015.6358> PMID: [26103030](https://pubmed.ncbi.nlm.nih.gov/26103030/).
  17. Wang T, Collet JP, Shapiro S, Ware MA. Adverse effects of medical cannabinoids: a systematic review. *Cmaj*. 2008; 178(13):1669–78. Epub 2008/06/19. <https://doi.org/10.1503/cmaj.071178> PMID: [18559804](https://pubmed.ncbi.nlm.nih.gov/18559804/); PubMed Central PMCID: PMC2413308.
  18. Velayudhan L, Van Diepen E, Marudkar M, Hands O, Suribhatla S, Prettyman R, et al. Therapeutic potential of cannabinoids in neurodegenerative disorders: a selective review. *Curr Pharm Des*. 2014; 20(13):2218–30. Epub 2013/07/09. <https://doi.org/10.2174/13816128113199990434> PMID: [23829360](https://pubmed.ncbi.nlm.nih.gov/23829360/).
  19. Beedham W, Sbai M, Allison I, Coary R, Shipway D. Cannabinoids in the Older Person: A Literature Review. *Geriatrics (Basel)*. 2020; 5(1). Epub 2020/01/17. <https://doi.org/10.3390/geriatrics5010002> PMID: [31941020](https://pubmed.ncbi.nlm.nih.gov/31941020/); PubMed Central PMCID: PMC7151062.
  20. Beauchet O. Medical cannabis use in older patients: Update on medical knowledge. *Maturitas*. 2018; 118:56–9. Epub 2018/11/13. <https://doi.org/10.1016/j.maturitas.2018.10.010> PMID: [30415756](https://pubmed.ncbi.nlm.nih.gov/30415756/).
  21. Abuhasira R, Schleider LB, Mechoulam R, Novack V. Epidemiological characteristics, safety and efficacy of medical cannabis in the elderly. *Eur J Intern Med*. 2018; 49:44–50. Epub 2018/02/06. <https://doi.org/10.1016/j.ejim.2018.01.019> PMID: [29398248](https://pubmed.ncbi.nlm.nih.gov/29398248/).
  22. O'Neill A, Wilson R, Blest-Hopley G, Annibale L, Colizzi M, Brammer M, et al. Normalization of mediotemporal and prefrontal activity, and mediotemporal-striatal connectivity, may underlie antipsychotic

- effects of cannabidiol in psychosis. *Psychol Med.* 2020;1–11. Epub 2020/01/30. <https://doi.org/10.1017/s0033291719003519> PMID: 31994476.
23. Crippa JA, Derenusson GN, Ferrari TB, Wichert-Ana L, Duran FL, Martin-Santos R, et al. Neural basis of anxiolytic effects of cannabidiol (CBD) in generalized social anxiety disorder: a preliminary report. *J Psychopharmacol.* 2011; 25(1):121–30. Epub 2010/09/11. <https://doi.org/10.1177/0269881110379283> PMID: 20829306.
  24. Appiah-Kusi E, Petros N, Wilson R, Colizzi M, Bossong MG, Valmaggia L, et al. Effects of short-term cannabidiol treatment on response to social stress in subjects at clinical high risk of developing psychosis. *Psychopharmacology (Berl).* 2020; 237(4):1121–30. Epub 2020/01/10. <https://doi.org/10.1007/s00213-019-05442-6> PMID: 31915861; PubMed Central PMCID: PMC7113209.
  25. Jones RT. Cardiovascular system effects of marijuana. *J Clin Pharmacol.* 2002; 42(S1):58s–63s. Epub 2002/11/05. <https://doi.org/10.1002/j.1552-4604.2002.tb06004.x> PMID: 12412837.
  26. Jadoon KA, Tan GD, O'Sullivan SE. A single dose of cannabidiol reduces blood pressure in healthy volunteers in a randomized crossover study. *JCI insight.* 2017; 2(12):e93760. <https://doi.org/10.1172/jci.insight.93760> PMID: 28614793.
  27. Englund A, Morrison PD, Nottage J, Hague D, Kane F, Bonaccorso S, et al. Cannabidiol inhibits THC-elicited paranoid symptoms and hippocampal-dependent memory impairment. *J Psychopharmacol.* 2013; 27(1):19–27. Epub 2012/10/09. <https://doi.org/10.1177/0269881112460109> PMID: 23042808.
  28. Hindocha C, Freeman TP, Schafer G, Gardener C, Das RK, Morgan CJ, et al. Acute effects of delta-9-tetrahydrocannabinol, cannabidiol and their combination on facial emotion recognition: a randomised, double-blind, placebo-controlled study in cannabis users. *Eur Neuropsychopharmacol.* 2015; 25(3):325–34. Epub 2014/12/24. <https://doi.org/10.1016/j.euroneuro.2014.11.014> PMID: 25534187; PubMed Central PMCID: PMC4398332.
  29. Moher D, Liberati A, Tetzlaff J, Altman DG, The PG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med.* 2009; 6(7):e1000097. <https://doi.org/10.1371/journal.pmed.1000097> PMID: 19621072
  30. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *Bmj.* 2008; 336(7650):924–6. Epub 2008/04/26. <https://doi.org/10.1136/bmj.39489.470347.AD> PMID: 18436948; PubMed Central PMCID: PMC2335261.
  31. Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol.* 2011; 64(4):401–6. Epub 2011/01/07. <https://doi.org/10.1016/j.jclinepi.2010.07.015> PMID: 21208779.
  32. Elbourne DR, Altman DG, Higgins JP, Curtin F, Worthington HV, Vail A. Meta-analyses involving crossover trials: methodological issues. *International Journal of Epidemiology.* 2002; 31(1):140–9. <https://doi.org/10.1093/ije/31.1.140> *International Journal of Epidemiology.* PMID: 11914310
  33. Tipton E, Pustejovsky JE. Small-Sample Adjustments for Tests of Moderators and Model Fit Using Robust Variance Estimation in Meta-Regression. *Journal of Educational and Behavioral Statistics.* 2015; 40(6):604–34. <https://doi.org/10.3102/1076998615606099>
  34. Pustejovsky JE, Tipton E. Small-Sample Methods for Cluster-Robust Variance Estimation and Hypothesis Testing in Fixed Effects Models. *Journal of Business & Economic Statistics.* 2018; 36(4):672–83. <https://doi.org/10.1080/07350015.2016.1247004>
  35. McCaffrey DF, Bell RM. Improved hypothesis testing for coefficients in generalized estimating equations with small samples of clusters. *Stat Med.* 2006; 25(23):4081–98. Epub 2006/02/04. <https://doi.org/10.1002/sim.2502> PMID: 16456895.
  36. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Bmj.* 1997; 315(7109):629–34. Epub 1997/10/06. <https://doi.org/10.1136/bmj.315.7109.629> PMID: 9310563; PubMed Central PMCID: PMC2127453.
  37. Duval S, Tweedie R. A Nonparametric “Trim and Fill” Method of Accounting for Publication Bias in Meta-Analysis. *Journal of the American Statistical Association.* 2000; 95(449):89–98. <https://doi.org/10.1080/01621459.2000.10473905>
  38. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package *Journal of Statistical Software.* 2010; 36(3):48.
  39. Volicer L, Stelly M, Morris J, McLaughlin J, Volicer BJ. Effects of dronabinol on anorexia and disturbed behavior in patients with Alzheimer's disease. *Int J Geriatr Psychiatry.* 1997; 12(9):913–9. Epub 1997/10/06. PMID: 9309469.
  40. Ahmed AI, van den Elsen GA, Colbers A, van der Marck MA, Burger DM, Feuth TB, et al. Safety and pharmacokinetics of oral delta-9-tetrahydrocannabinol in healthy older subjects: a randomized controlled trial. *Eur Neuropsychopharmacol.* 2014; 24(9):1475–82. Epub 2014/07/19. <https://doi.org/10.1016/j.euroneuro.2014.06.007> PMID: 25035121.

41. Walther S, Schupbach B, Seifritz E, Homan P, Strik W. Randomized, controlled crossover trial of dronabinol, 2.5 mg, for agitation in 2 patients with dementia. *J Clin Psychopharmacol*. 2011; 31(2):256–8. Epub 2011/03/03. <https://doi.org/10.1097/JCP.0b013e31820e861c> PMID: [21364345](#).
42. Pickering EE, Semple SJ, Nazir MS, Murphy K, Snow TM, Cummin AR, et al. Cannabinoid effects on ventilation and breathlessness: a pilot study of efficacy and safety. *Chron Respir Dis*. 2011; 8(2):109–18. Epub 2011/03/26. <https://doi.org/10.1177/1479972310391283> PMID: [21436223](#).
43. Lane M, Vogel CL, Ferguson J, Krasnow S, Saiers JL, Hamm J, et al. Dronabinol and prochlorperazine in combination for treatment of cancer chemotherapy-induced nausea and vomiting. *J Pain Symptom Manage*. 1991; 6(6):352–9. Epub 1991/08/01. [https://doi.org/10.1016/0885-3924\(91\)90026-z](https://doi.org/10.1016/0885-3924(91)90026-z) PMID: [1652611](#).
44. Sieradzan KA, Fox SH, Hill M, Dick JP, Crossman AR, Brotchie JM. Cannabinoids reduce levodopa-induced dyskinesia in Parkinson's disease: a pilot study. *Neurology*. 2001; 57(11):2108–11. Epub 2001/12/12. <https://doi.org/10.1212/wnl.57.11.2108> PMID: [11739835](#).
45. Jatoi A, Windschitl HE, Loprinzi CL, Sloan JA, Dakhil SR, Mailliard JA, et al. Dronabinol versus megestrol acetate versus combination therapy for cancer-associated anorexia: a North Central Cancer Treatment Group study. *J Clin Oncol*. 2002; 20(2):567–73. Epub 2002/01/12. <https://doi.org/10.1200/JCO.2002.20.2.567> PMID: [11786587](#).
46. Zajicek J, Fox P, Sanders H, Wright D, Vickery J, Nunn A, et al. Cannabinoids for treatment of spasticity and other symptoms related to multiple sclerosis (CAMS study): multicentre randomised placebo-controlled trial. *The Lancet*. 2003; 362(9395):1517–26. [https://doi.org/10.1016/S0140-6736\(03\)14738-1](https://doi.org/10.1016/S0140-6736(03)14738-1) PMID: [14615106](#)
47. Svendsen KB, Jensen TS, Bach FW. Does the cannabinoid dronabinol reduce central pain in multiple sclerosis? Randomised double blind placebo controlled crossover trial. *Bmj*. 2004; 329(7460):253. Epub 2004/07/20. <https://doi.org/10.1136/bmj.38149.566979.AE> PMID: [15258006](#); PubMed Central PMCID: PMC498019.
48. Zajicek JP, Sanders HP, Wright DE, Vickery PJ, Ingram WM, Reilly SM, et al. Cannabinoids in multiple sclerosis (CAMS) study: safety and efficacy data for 12 months follow up. *J Neurol Neurosurg Psychiatry*. 2005; 76(12):1664–9. Epub 2005/11/18. <https://doi.org/10.1136/jnnp.2005.070136> PMID: [16291891](#); PubMed Central PMCID: PMC1739436.
49. Strasser F, Luftner D, Possinger K, Ernst G, Ruhstaller T, Meissner W, et al. Comparison of orally administered cannabis extract and delta-9-tetrahydrocannabinol in treating patients with cancer-related anorexia-cachexia syndrome: a multicenter, phase III, randomized, double-blind, placebo-controlled clinical trial from the Cannabis-In-Cachexia-Study-Group. *J Clin Oncol*. 2006; 24(21):3394–400. Epub 2006/07/20. <https://doi.org/10.1200/JCO.2005.05.1847> PMID: [16849753](#).
50. Tomida I, Azuara-Blanco A, House H, Flint M, Pertwee RG, Robson PJ. Effect of sublingual application of cannabinoids on intraocular pressure: a pilot study. *J Glaucoma*. 2006; 15(5):349–53. Epub 2006/09/22. <https://doi.org/10.1097/01.jig.0000212260.04488.60> PMID: [16988594](#).
51. Meiri E, Jhangiani H, Vredenburg JJ, Barbato LM, Carter FJ, Yang HM, et al. Efficacy of dronabinol alone and in combination with ondansetron versus ondansetron alone for delayed chemotherapy-induced nausea and vomiting. *Curr Med Res Opin*. 2007; 23(3):533–43. Epub 2007/03/16. <https://doi.org/10.1185/030079907x167525> PMID: [17355735](#).
52. Curtis A, Mitchell I, Patel S, Ives N, Rickards H. A pilot study using nabilone for symptomatic treatment in Huntington's disease. *Mov Disord*. 2009; 24(15):2254–9. Epub 2009/10/22. <https://doi.org/10.1002/mds.22809> PMID: [19845035](#).
53. Johnson JR, Burnell-Nugent M, Lossignol D, Ganae-Motan ED, Potts R, Fallon MT. Multicenter, double-blind, randomized, placebo-controlled, parallel-group study of the efficacy, safety, and tolerability of THC:CBD extract and THC extract in patients with intractable cancer-related pain. *J Pain Symptom Manage*. 2010; 39(2):167–79. Epub 2009/11/10. <https://doi.org/10.1016/j.jpainsymman.2009.06.008> PMID: [19896326](#).
54. Ware MA, Fitzcharles MA, Joseph L, Shir Y. The effects of nabilone on sleep in fibromyalgia: results of a randomized controlled trial. *Anesth Analg*. 2010; 110(2):604–10. Epub 2009/12/17. <https://doi.org/10.1213/ANE.0b013e3181c76f70> PMID: [20007734](#).
55. Weber M, Goldman B, Truniger S. Tetrahydrocannabinol (THC) for cramps in amyotrophic lateral sclerosis: a randomised, double-blind crossover trial. *J Neurol Neurosurg Psychiatry*. 2010; 81(10):1135–40. Epub 2010/05/26. <https://doi.org/10.1136/jnnp.2009.200642> PMID: [20498181](#).
56. Brisbois TD, de Kock IH, Watanabe SM, Mirhosseini M, Lamoureux DC, Chasen M, et al. Delta-9-tetrahydrocannabinol may palliate altered chemosensory perception in cancer patients: results of a randomized, double-blind, placebo-controlled pilot trial. *Ann Oncol*. 2011; 22(9):2086–93. Epub 2011/02/24. <https://doi.org/10.1093/annonc/mdq727> PMID: [21343383](#).



57. Zadikoff C, Wadia PM, Miyasaki J, Chen R, Lang AE, So J, et al. Cannabinoid, CB1 agonists in cervical dystonia: Failure in a phase IIa randomized controlled trial. *Basal Ganglia*. 2011; 1(2):91–5. <https://doi.org/10.1016/j.baga.2011.04.002>.
58. Toth C, Mawani S, Brady S, Chan C, Liu C, Mehina E, et al. An enriched-enrolment, randomized withdrawal, flexible-dose, double-blind, placebo-controlled, parallel assignment efficacy study of nabilone as adjuvant in the treatment of diabetic peripheral neuropathic pain. *Pain*. 2012; 153(10):2073–82. Epub 2012/08/28. <https://doi.org/10.1016/j.pain.2012.06.024> PMID: 22921260.
59. Zajicek J, Ball S, Wright D, Vickery J, Nunn A, Miller D, et al. Effect of dronabinol on progression in progressive multiple sclerosis (CUPID): a randomised, placebo-controlled trial. *Lancet Neurol*. 2013; 12(9):857–65. Epub 2013/07/17. [https://doi.org/10.1016/S1474-4422\(13\)70159-5](https://doi.org/10.1016/S1474-4422(13)70159-5) PMID: 23856559; PubMed Central PMCID: PMC3744749.
60. Ahmed AI, van den Elsen GA, Colbers A, Kramers C, Burger DM, van der Marck MA, et al. Safety, pharmacodynamics, and pharmacokinetics of multiple oral doses of delta-9-tetrahydrocannabinol in older persons with dementia. *Psychopharmacology (Berl)*. 2015; 232(14):2587–95. Epub 2015/03/11. <https://doi.org/10.1007/s00213-015-3889-y> PMID: 25752889; PubMed Central PMCID: PMC4480847.
61. van den Elsen GA, Ahmed AI, Verkes RJ, Kramers C, Feuth T, Rosenberg PB, et al. Tetrahydrocannabinol for neuropsychiatric symptoms in dementia: A randomized controlled trial. *Neurology*. 2015; 84(23):2338–46. Epub 2015/05/15. <https://doi.org/10.1212/WNL.0000000000001675> PMID: 25972490; PubMed Central PMCID: PMC4464746.
62. de Vries M, Van Rijckevorsel DC, Vissers KC, Wilder-Smith OH, Van Goor H. Single dose delta-9-tetrahydrocannabinol in chronic pancreatitis patients: analgesic efficacy, pharmacokinetics and tolerability. *Br J Clin Pharmacol*. 2016; 81(3):525–37. Epub 2015/10/28. <https://doi.org/10.1111/bcp.12811> PMID: 26505163; PubMed Central PMCID: PMC4767190.
63. van Amerongen G, Kanhai K, Baakman AC, Heuberger J, Klaassen E, Beumer TL, et al. Effects on Spasticity and Neuropathic Pain of an Oral Formulation of Delta9-tetrahydrocannabinol in Patients With Progressive Multiple Sclerosis. *Clin Ther*. 2018; 40(9):1467–82. Epub 2017/02/13. <https://doi.org/10.1016/j.clinthera.2017.01.016> PMID: 28189366.
64. Carley DW, Prasad B, Reid KJ, Malkani R, Attarian H, Abbott SM, et al. Pharmacotherapy of Apnea by Cannabimimetic Enhancement, the PACE Clinical Trial: Effects of Dronabinol in Obstructive Sleep Apnea. *Sleep*. 2018; 41(1). <https://doi.org/10.1093/sleep/zsx184> PMID: 29121334; PubMed Central PMCID: PMC5806568.
65. Herrmann N, Ruthirakuhan M, Gallagher D, Verhoeff N, Kiss A, Black SE, et al. Randomized Placebo-Controlled Trial of Nabilone for Agitation in Alzheimer's Disease. *Am J Geriatr Psychiatry*. 2019; 27(11):1161–73. Epub 2019/06/12. <https://doi.org/10.1016/j.jagp.2019.05.002> PMID: 31182351.
66. van den Elsen GAH, Ahmed AIA, Verkes RJ, Feuth T, van der Marck MA, Olde Rikkert MGM. Tetrahydrocannabinol in Behavioral Disturbances in Dementia: A Crossover Randomized Controlled Trial. *Am J Geriatr Psychiatry*. 2015; 23(12):1214–24. Epub 2015/11/13. <https://doi.org/10.1016/j.jagp.2015.07.011> PMID: 26560511.
67. Peball M, Krismer F, Knaus HG, Djamshidian A, Werkmann M, Carbone F, et al. Non-Motor Symptoms in Parkinson's Disease are Reduced by Nabilone. *Ann Neurol*. 2020; 88(4):712–. Epub 2020/08/07. <https://doi.org/10.1002/ana.25864> PMID: 32757413; PubMed Central PMCID: PMC7540547.
68. Carroll CB, Bain P, Teare L, Liu X, Joint C, Wroath C, et al. Cannabis for dyskinesia in Parkinson disease: A randomized double-blind crossover study. *Neurology*. 2004; 63(7):1245–50. 2004-19648-015. <https://doi.org/10.1212/01.wnl.0000140288.48796.8e> PMID: 15477546
69. Vaney C, Heinzl-Gutenbrunner M, Jobin P, Tschopp F, Gattlen B, Hagen U, et al. Efficacy, safety and tolerability of an orally administered cannabis extract in the treatment of spasticity in patients with multiple sclerosis: a randomized, double-blind, placebo-controlled, crossover study. *Mult Scler*. 2004; 10(4):417–24. Epub 2004/08/26. <https://doi.org/10.1191/1352458504ms10480a> PMID: 15327040.
70. Wade DT, Makela P, Robson P, House H, Bateman C. Do cannabis-based medicinal extracts have general or specific effects on symptoms in multiple sclerosis? A double-blind, randomized, placebo-controlled study on 160 patients. *Mult Scler*. 2004; 10(4):434–41. Epub 2004/08/26. <https://doi.org/10.1191/1352458504ms10820a> PMID: 15327042.
71. Blake DR, Robson P, Ho M, Jubb RW, McCabe CS. Preliminary assessment of the efficacy, tolerability and safety of a cannabis-based medicine (Sativex) in the treatment of pain caused by rheumatoid arthritis. *Rheumatology (Oxford)*. 2006; 45(1):50–2. Epub 2005/11/12. <https://doi.org/10.1093/rheumatology/kei183> PMID: 16282192.
72. Nurmikko TJ, Serpell MG, Hoggart B, Toomey PJ, Morlion BJ, Haines D. Sativex successfully treats neuropathic pain characterised by allodynia: a randomised, double-blind, placebo-controlled clinical trial. *Pain*. 2007; 133(1–3):210–20. Epub 2007/11/13. <https://doi.org/10.1016/j.pain.2007.08.028> PMID: 17997224.



73. Duran M, Perez E, Abanades S, Vidal X, Saura C, Majem M, et al. Preliminary efficacy and safety of an oromucosal standardized cannabis extract in chemotherapy-induced nausea and vomiting. *Br J Clin Pharmacol*. 2010; 70(5):656–63. Epub 2010/11/03. <https://doi.org/10.1111/j.1365-2125.2010.03743.x> PMID: [21039759](https://pubmed.ncbi.nlm.nih.gov/21039759/); PubMed Central PMCID: PMC2997305.
74. Johnson JR, Burnell-Nugent M, Lossignol D, Ganae-Motan ED, Potts R, Fallon MT. Multicenter, double-blind, randomized, placebo-controlled, parallel-group study of the efficacy, safety, and tolerability of THC:CBD extract and THC extract in patients with intractable cancer-related pain. *Journal of Pain & Symptom Management*. 2010; 39(2):167–79. <https://doi.org/10.1016/j.jpainsymman.2009.06.008> PMID: [105129735](https://pubmed.ncbi.nlm.nih.gov/105129735/). Language: English. Entry Date: 20100521. Revision Date: 20150711. Publication Type: Journal Article.
75. Notcutt W, Langford R, Davies P, Ratcliffe S, Potts R. A placebo-controlled, parallel-group, randomized withdrawal study of subjects with symptoms of spasticity due to multiple sclerosis who are receiving long-term Sativex(R) (nabiximols). *Mult Scler*. 2012; 18(2):219–28. Epub 2011/09/01. <https://doi.org/10.1177/1352458511419700> PMID: [21878454](https://pubmed.ncbi.nlm.nih.gov/21878454/).
76. Portenoy RK, Ganae-Motan ED, Allende S, Yanagihara R, Shaiova L, Weinstein S, et al. Nabiximols for opioid-treated cancer patients with poorly-controlled chronic pain: a randomized, placebo-controlled, graded-dose trial. *J Pain*. 2012; 13(5):438–49. Epub 2012/04/10. <https://doi.org/10.1016/j.jpain.2012.01.003> PMID: [22483680](https://pubmed.ncbi.nlm.nih.gov/22483680/).
77. Zajicek JP, Hobart JC, Slade A, Barnes D, Mattison PG. Multiple sclerosis and extract of cannabis: results of the MUSEC trial. *J Neurol Neurosurg Psychiatry*. 2012; 83(11):1125–32. Epub 2012/07/14. <https://doi.org/10.1136/jnnp-2012-302468> PMID: [22791906](https://pubmed.ncbi.nlm.nih.gov/22791906/).
78. Lynch ME, Cesar-Rittenberg P, Hohmann AG. A double-blind, placebo-controlled, crossover pilot trial with extension using an oral mucosal cannabinoid extract for treatment of chemotherapy-induced neuropathic pain. *J Pain Symptom Manage*. 2014; 47(1):166–73. Epub 2013/06/08. <https://doi.org/10.1016/j.jpainsymman.2013.02.018> PMID: [23742737](https://pubmed.ncbi.nlm.nih.gov/23742737/).
79. Serpell M, Ratcliffe S, Hovorka J, Schofield M, Taylor L, Lauder H, et al. A double-blind, randomized, placebo-controlled, parallel group study of THC/CBD spray in peripheral neuropathic pain treatment. *Eur J Pain*. 2014; 18(7):999–1012. Epub 2014/01/15. <https://doi.org/10.1002/j.1532-2149.2013.00445.x> PMID: [24420962](https://pubmed.ncbi.nlm.nih.gov/24420962/).
80. Jadoon KA, Ratcliffe SH, Barrett DA, Thomas EL, Stott C, Bell JD, et al. Efficacy and Safety of Cannabidiol and Tetrahydrocannabinol on Glycemic and Lipid Parameters in Patients With Type 2 Diabetes: A Randomized, Double-Blind, Placebo-Controlled, Parallel Group Pilot Study. *Diabetes Care*. 2016; 39(10):1777–86. Epub 2016/08/31. <https://doi.org/10.2337/dc16-0650> PMID: [27573936](https://pubmed.ncbi.nlm.nih.gov/27573936/).
81. Fallon MT, Albert Lux E, McQuade R, Rossetti S, Sanchez R, Sun W, et al. Sativex oromucosal spray as adjunctive therapy in advanced cancer patients with chronic pain unrelieved by optimized opioid therapy: two double-blind, randomized, placebo-controlled phase 3 studies. *British journal of pain*. 2017; 11(3):119–33. Epub 05/17. <https://doi.org/10.1177/2049463717710042> PMID: [28785408](https://pubmed.ncbi.nlm.nih.gov/28785408/).
82. Lichtman AH, Lux EA, McQuade R, Rossetti S, Sanchez R, Sun W, et al. Results of a Double-Blind, Randomized, Placebo-Controlled Study of Nabiximols Oromucosal Spray as an Adjunctive Therapy in Advanced Cancer Patients with Chronic Uncontrolled Pain. *Journal of Pain and Symptom Management*. 2018; 55(2):179–88.e1. <https://doi.org/10.1016/j.jpainsymman.2017.09.001> PMID: [28923526](https://pubmed.ncbi.nlm.nih.gov/28923526/)
83. Marková J, Essner U, Akmaz B, Marinelli M, Trompke C, Lentschat A, et al. Sativex® as add-on therapy vs. further optimized first-line ANTispastics (SAVANT) in resistant multiple sclerosis spasticity: a double-blind, placebo-controlled randomised clinical trial. *International Journal of Neuroscience*. 2019; 129(2):119–28. <https://doi.org/10.1080/00207454.2018.1481066> PMID: [29792372](https://pubmed.ncbi.nlm.nih.gov/29792372/)
84. Riva N, Mora G, Sorarù G, Lunetta C, Ferraro OE, Falzone Y, et al. Safety and efficacy of nabiximols on spasticity symptoms in patients with motor neuron disease (CANALS): a multicentre, double-blind, randomised, placebo-controlled, phase 2 trial. *Lancet Neurol*. 2019; 18(2):155–64. Epub 2018/12/18. [https://doi.org/10.1016/S1474-4422\(18\)30406-X](https://doi.org/10.1016/S1474-4422(18)30406-X) PMID: [30554828](https://pubmed.ncbi.nlm.nih.gov/30554828/).
85. Consroe P, Laguna J, Allender J, Snider S, Stern L, Sandyk R, et al. Controlled clinical trial of cannabidiol in Huntington's disease. *Pharmacol Biochem Behav*. 1991; 40(3):701–8. Epub 1991/11/01. [https://doi.org/10.1016/0091-3057\(91\)90386-g](https://doi.org/10.1016/0091-3057(91)90386-g) PMID: [1839644](https://pubmed.ncbi.nlm.nih.gov/1839644/).
86. Iffland K, Grotenhermen F. An Update on Safety and Side Effects of Cannabidiol: A Review of Clinical Data and Relevant Animal Studies. *Cannabis Cannabinoid Res*. 2017; 2(1):139–54. Epub 2017/09/02. <https://doi.org/10.1089/can.2016.0034> PMID: [28861514](https://pubmed.ncbi.nlm.nih.gov/28861514/); PubMed Central PMCID: PMC5569602.
87. Black N, Stockings E, Campbell G, Tran LT, Zagic D, Hall WD, et al. Cannabinoids for the treatment of mental disorders and symptoms of mental disorders: a systematic review and meta-analysis. *Lancet Psychiatry*. 2019; 6(12):995–1010. Epub 2019/11/02. [https://doi.org/10.1016/S2215-0366\(19\)30401-8](https://doi.org/10.1016/S2215-0366(19)30401-8) PMID: [31672337](https://pubmed.ncbi.nlm.nih.gov/31672337/); PubMed Central PMCID: PMC6949116.
88. Bueno-Cavanillas A, Padilla-Ruiz F, Jiménez-Moleón JJ, Peinado-Alonso CA, Gálvez-Vargas R. Risk factors in falls among the elderly according to extrinsic and intrinsic precipitating causes. *Eur J*

- Epidemiol. 2000; 16(9):849–59. Epub 2001/04/12. <https://doi.org/10.1023/a:1007636531965> PMID: [11297228](https://pubmed.ncbi.nlm.nih.gov/11297228/).
89. Khow KSF, Visvanathan R. Falls in the Aging Population. *Clin Geriatr Med*. 2017; 33(3):357–68. Epub 2017/07/12. <https://doi.org/10.1016/j.cger.2017.03.002> PMID: [28689568](https://pubmed.ncbi.nlm.nih.gov/28689568/).
  90. Abuhasira R, Ron A, Sikorin I, Novack V. Medical Cannabis for Older Patients-Treatment Protocol and Initial Results. *J Clin Med*. 2019; 8(11). Epub 2019/11/07. <https://doi.org/10.3390/jcm8111819> PMID: [31683817](https://pubmed.ncbi.nlm.nih.gov/31683817/); PubMed Central PMCID: PMC6912698.
  91. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials*. 1996; 17(1):1–12. Epub 1996/02/01. [https://doi.org/10.1016/0197-2456\(95\)00134-4](https://doi.org/10.1016/0197-2456(95)00134-4) PMID: [8721797](https://pubmed.ncbi.nlm.nih.gov/8721797/).
  92. Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet*. 2012; 380(9836):37–43. Epub 2012/05/15. [https://doi.org/10.1016/S0140-6736\(12\)60240-2](https://doi.org/10.1016/S0140-6736(12)60240-2) PMID: [22579043](https://pubmed.ncbi.nlm.nih.gov/22579043/).
  93. Wastesson JW, Morin L, Tan ECK, Johnell K. An update on the clinical consequences of polypharmacy in older adults: a narrative review. *Expert Opin Drug Saf*. 2018; 17(12):1185–96. Epub 2018/12/13. <https://doi.org/10.1080/14740338.2018.1546841> PMID: [30540223](https://pubmed.ncbi.nlm.nih.gov/30540223/).
  94. Imrey PB. Limitations of Meta-analyses of Studies With High Heterogeneity. *JAMA Netw Open*. 2020; 3(1):e1919325. Epub 2020/01/11. <https://doi.org/10.1001/jamanetworkopen.2019.19325> PMID: [31922554](https://pubmed.ncbi.nlm.nih.gov/31922554/).
  95. Singh S, Bajorek B. Pharmacotherapy in the ageing patient: The impact of age per se (A review). *Ageing Res Rev*. 2015; 24(Pt B):99–110. Epub 2015/08/01. <https://doi.org/10.1016/j.arr.2015.07.006> PMID: [26226330](https://pubmed.ncbi.nlm.nih.gov/26226330/).