

CORONAVIRUS

Cannabidiol inhibits SARS-CoV-2 replication through induction of the host ER stress and innate immune responses

Long Chi Nguyen¹§, Dongbo Yang¹§, Vlad Nicolaescu^{2,10}§, Thomas J. Best³§, Haley Gula^{2,10}, Divyasha Saxena⁴, Jon D. Gabbard⁴, Shao-Nong Chen⁵, Takashi Ohtsuki⁵†, John Brent Friesen⁵, Nir Drayman⁶, Adil Mohamed⁶, Christopher Dann¹, Diane Silva⁷, Lydia Robinson-Mailman¹, Andrea Valdespino¹, Letícia Stock¹, Eva Suárez¹, Krysten A. Jones⁸, Saara-Anne Azizi⁸, Jennifer K. Demarco⁴, William E. Severson⁴, Charles D. Anderson⁴, James Michael Millis⁹, Bryan C. Dickinson⁸, Savaş Tay⁶, Scott A. Oakes⁷, Guido F. Pauli⁵, Kenneth E. Palmer⁴, The National COVID Cohort Collaborative Consortium‡, David O. Meltzer³, Glenn Randall^{2,10}*, Marsha Rich Rosner¹*

The spread of SARS-CoV-2 and ongoing COVID-19 pandemic underscores the need for new treatments. Here we report that cannabidiol (CBD) inhibits infection of SARS-CoV-2 in cells and mice. CBD and its metabolite 7-OH-CBD, but not THC or other congeneric cannabinoids tested, potently block SARS-CoV-2 replication in lung epithelial cells. CBD acts after viral entry, inhibiting viral gene expression and reversing many effects of SARS-CoV-2 on host gene transcription. CBD inhibits SARS-CoV-2 replication in part by up-regulating the host IRE1 α RNase endoplasmic reticulum (ER) stress response and interferon signaling pathways. In matched groups of human patients from the National COVID Cohort Collaborative, CBD (100 mg/ml oral solution per medical records) had a significant negative association with positive SARS-CoV-2 tests. This study highlights CBD as a potential preventative agent for early-stage SARS-CoV-2 infection and merits future clinical trials. We caution against use of non-medical formulations including edibles, inhalants or topicals as a preventative or treatment therapy at the present time.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is responsible for coronavirus disease 2019 (COVID-19), a pandemic that continues to cause widespread morbidity and mortality across the globe. SARS-CoV-2 is the seventh species of coronavirus known to infect people. These coronaviruses, which include SARS-CoV, 229E, NL63, OC43, HKU1, and MERS-CoV, cause a range of symptoms from the common cold to more severe pathologies (1). Despite recent vaccine availability, SARS-CoV-2 is still spreading rapidly (2), highlighting the need for alternative treatments, especially for populations with limited inclination or access to vaccines. To date, few therapies have been identified that block SARS-CoV-2 replication and viral production.

SARS-CoV-2 is a positive-sense single-stranded RNA (+ssRNA) enveloped virus composed of a lipid bilayer and four structural

proteins that drive viral particle formation. The spike (S), membrane (M), and envelope (E) are integral proteins of the virus membrane and promote virion budding while also recruiting the nucleocapsid (N) protein and the viral genomic RNA into nascent virions. Like its close relative SARS-CoV, SARS-CoV-2 primarily enters human cells by the binding of the viral S protein to the angiotensin converting enzyme 2 (ACE2) receptor (3–5), after which the S protein undergoes proteolysis by transmembrane protease serine 2 (TMPRSS2) or other proteases into two non-covalently bound peptides (S1, S2) that facilitate viral entry into the host cell. The N-terminal S1 binds the ACE2 receptor, and the C-terminal S2 mediates viral-cell membrane fusion following proteolytic cleavage. Depending upon the cell type, viral entry can also occur after ACE2 binding, independent of proteolytic cleavage (6–8). Following cell entry, the SARS-CoV-2 genome is translated into two large polypeptides that are cleaved by two viral proteases, Mpro and PLpro (9, 10), to produce 15 proteins, in addition to the synthesis of subgenomic RNAs that encode another 10 accessory proteins plus the 4 structural proteins. These proteins enable viral replication, assembly, and budding. In an effort to suppress infection by the SARS-CoV-2 beta-coronavirus as well as other evolving pathogenic viruses, we tested the antiviral potential of a number of small molecules that target host stress response pathways.

One potential regulator of the host stress and antiviral inflammatory responses is cannabidiol (CBD), a member of the cannabinoid class of natural products (11) produced by *Cannabis sativa* (Cannabaceae; marijuana/hemp). Hemp refers to cannabis plants or materials derived thereof that contain 0.3% or less of the psychotropic tetrahydrocannabinol (THC) and typically have relatively high CBD content. By contrast, marijuana refers to *C. sativa* materials with more than 0.3% THC by dry weight. THC acts through binding to

¹Ben May Department for Cancer Research, University of Chicago, Chicago, IL 60637. ²Department of Microbiology, University of Chicago, Chicago, IL 60637. ³Center for Health and the Social Sciences, University of Chicago, Chicago IL 60637. ⁴Center for Predictive Medicine for Biodefense and Emerging Infectious Diseases, University of Louisville, Louisville KY 40222. ⁵Pharmacognosy Institute and Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois at Chicago, Chicago, IL 60612. ⁶Pritzker School of Molecular Engineering, University of Chicago, Chicago IL 60637. ⁷Department of Pathology, University of Chicago, Chicago IL 60637. ⁸Department of Chemistry, University of Chicago, Chicago IL 60637. ⁹Department of Surgery, University of Chicago, Chicago IL 60637. ¹⁰Howard Taylor Ricketts Laboratory, Argonne National Laboratory, Lemont IL 60439.

*Corresponding author. Email: grandall@bsd.uchicago.edu (G.R.); mrosner@uchicago.edu (M.R.R.)

†Present address: Department of Food Bioscience and Biotechnology, College of Bioresource Sciences, Nihon University, Fujisawa, Kanagawa, Japan.

‡Consortium authors and affiliations are provided in the Supplementary Materials. §Equal contribution.

MS no:	Emp name:	Date / Time:	PE's:	AA's:	Comments:	Art no:
RAabi6110/IMMUNOLOGY, LIFE SCIENCES	jrnindesign	1-20-2022 / 22:52				

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

the cannabinoid receptor, and CBD potentiates this interaction (11). Despite numerous studies and many unsubstantiated claims related to CBD-containing products, the biologic actions of CBD itself are unclear and specific targets are mostly unknown (12). However, an oral solution of CBD is an FDA-approved drug, largely for the treatment of epilepsy (13). Thus, CBD has drug status, is viable as a therapeutic, and cannot be marketed as a dietary supplement in the United States (12). Although limited, some studies have reported that certain cannabinoids have antiviral effects against hepatitis C virus (HCV) and other viruses (14).

RESULTS

High purity CBD inhibits SARS-CoV-2 replication in human lung epithelial cells

To test the effect of CBD on SARS-CoV-2 replication, we pretreated A549 human lung carcinoma cells expressing exogenous human ACE-2 receptor (A549-ACE2) for 2 hours with 0–10 μM CBD prior to infection with SARS-CoV-2. After 48 hours, we monitored cells for expression of the viral spike protein (S) and viral titer. CBD potently inhibited viral replication under non-toxic conditions with an EC₅₀ of ~ 1 μM (Fig. 1A; fig. S1A). CBD inhibited SARS-CoV-2 replication in human Calu3 lung and Vero E6 monkey kidney epithelial cells as well (fig. S1B), and no toxicity was observed at the effective doses (fig. S1C,D). Finally, we tested three SARS-CoV-2 variants of concern (α , β , and γ) in addition to the original SARS-CoV-2 strain, and their ability to infect cells was comparably inhibited by CBD (Fig. 1C).

When isolated from its source plant, natural non-synthetic CBD is typically extracted along with other cannabinoids, representing

the unavoidable residual complexity of natural products (12). To verify that CBD is indeed responsible for the viral inhibition, we analyzed a CBD reference standard as well as CBD from four different sources for purity using 100% quantitative NMR (qNMR). These sources included two chemical vendors (Suppliers A and B) and two commercial vendors (Suppliers C and D). The striking congruence between the experimental ¹H NMR and the recently established quantum-mechanical HiFSA (¹H Iterative Full Spin Analysis) profiles observed for all materials confirmed that 1) the compounds used were indeed CBD with purities of at least 97% (Fig. 1B) and 2) congeneric cannabinoids were not present at levels above 1.0%. Analysis of these different CBD samples in the viral A549-ACE2 infection assay showed similar EC₅₀s with a range from 0.6–1.8 μM , likely reflecting the intrinsic variability of the biological assay (Fig. 1A). No toxicity was observed for any of the CBD preparations at the doses used to inhibit viral infection (fig. S1 E-G).

The CBD metabolite 7-OH-CBD, but not a panel of closely related CBD congeners, exhibits antiviral activity

CBD is often consumed as part of a *C. sativa* extract, particularly in combination with psychoactive THC enriched in marijuana plants. We therefore determined whether congeneric cannabinoids, especially analogues with closely related structures and polarities produced by the hemp plant, are also capable of inhibiting SARS-CoV-2 infection. Remarkably, of this group, only CBD was a potent agent, while no or very limited antiviral activity was exhibited by these structurally closely related congeners that share biosynthesis pathways and form the biogenetically determined residual complexity of CBD purified from *C. sativa*: THC, cannabidiolic acid (CBDA), cannabidivarin (CBDV), cannabichromene (CBC), or cannabigerol




Fig. 1. Cannabidiol (CBD) is a potent inhibitor of SARS-CoV-2 infection *in vitro*. (A) A549-ACE2 cells were treated with indicated doses of CBD from four different suppliers followed by infection with SARS-CoV-2 at an MOI of 0.5 for 48 hours. The cells were stained for spike protein and the percentage of cells expressing the spike protein in each condition was plotted. EC₅₀ values are indicated. (B) The ¹H qNMR spectra of CBD reference material and CBD samples from four different suppliers. (C) A549-ACE2 cells were treated with CBD from supplier A followed by infection with SARS-CoV-2 or α , β or γ variants at an MOI of 0.5 for 48 hours. The cells were stained for spike protein and the percentage of cells expressing the spike protein in each condition was plotted. EC₅₀ values are indicated.

MS no:	Emp name:	Date / Time:	PE's:	AA's:	Comments:	Art no:
RAabi6110/IMMUNOLOGY, LIFE SCIENCES	jrnindesign	1-20-2022 / 22:52				Fig 1

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

F2 (CBG) (Fig. 2 A,D; see Methods). None of these cannabinoids were toxic to the A549-ACE2 cells in the dose range of interest (fig. S2). Notably, combining CBD with THC (1:1) significantly suppressed CBD efficacy consistent with competitive inhibition by THC.

CBD is rapidly metabolized in the intestine and liver into two main metabolites, 7-carboxy-cannabidiol (7-COOH-CBD) and 7-hydroxy-cannabidiol (7-OH-CBD). The level of 7-COOH-CBD is 40-fold higher, and the level of 7-OH-CBD is 38% of the CBD level in human plasma (15). CBD and its 7-OH-CBD metabolite are the active and equipotent ingredients for the treatment of epilepsy (13). Like CBD, 7-OH-CBD effectively inhibited SARS-CoV-2 replication in A549-ACE2 cells (Fig. 2C) and was non-toxic to cells (fig. S2H,I). Analysis of blood plasma levels in healthy patients taking 1500 mg daily of FDA-approved CBD solution (Epidiolex) showed a maximal concentration (C_{max}) at 7 days for CBD and 7-OH-CBD of 1.7 μ M and 0.56 μ M, respectively; the C_{max} can be further increased several-fold by co-administration with a high-fat meal (15). Taken in aggregate, these results suggest the effective plasma concentrations of CBD and its metabolite are within the therapeutic range to inhibit SARS-CoV-2 infection in humans.

CBD acts at an early step after viral entry into cells

CBD could be acting by blocking viral entry to host cells or at later steps following infection. As CBD was reported to decrease ACE2

expression in some epithelial cells, including A549 (16), we first determined whether CBD suppressed the SARS-CoV-2 receptor in the A549-ACE2, Calu-3, and Vero E6 cells. No decrease in ACE2 expression was observed (Fig. 3A; fig. S4A,B). Furthermore, analysis of lentiviruses pseudotyped with either the SARS-CoV-2 spike protein or the vesicular stomatitis virus (VSV) glycoprotein (17) showed that 10 μ M CBD only weakly inhibited cell entry by spike-expressing virus, suggesting that other mechanisms are largely responsible for its antiviral effects. The robustness of the assay was confirmed by using anti-spike antibodies that effectively blocked viral infection of lentivirus pseudotyped with spike, but not VSVg (Fig. 3B, and figs. S3 A and B). In contrast to the negligible effect on viral entry, CBD was very effective (~95–99%) at inhibiting SARS-CoV-2 spike protein expression in host cells at 2 and 6 hours after infection post entry (Fig. 3C). This was true even in the presence of antibodies to the spike protein to prevent reinfection (Fig. 3D) suggesting CBD acts early in the infection cycle, in a post entry step. CBD was also partially effective (~60%) at inhibiting SARS-CoV-2 at 15 h after infection (Fig. 3C), suggesting a possible secondary effect on viral assembly and release. To assess whether CBD might be preventing viral protein processing by the viral proteases Mpro or PLpro, we assayed their activity *in vitro* (fig. S4C,D). CBD did not affect the activity of either protease, raising the possibility that CBD targets host cell processes.




Fig. 2. Limited or no inhibition of SARS-CoV-2 infection by Cannabinoids other than CBD. (A) A549-ACE2 cells were treated with indicated doses of various cannabinoids or a CBD/THC 1:1 mixture followed by infection with SARS-CoV-2 at an MOI of 0.5 for 48 hours. The cells were stained for spike protein and the percentage of cells expressing the spike protein in each condition was plotted. All cannabinoids tested were isolated from a hemp extract as described in Methods. (B) Chemical structures of cannabinoids and 7-OH-CBD. (C) A549-ACE2 cells were treated with indicated doses of 7-OH-CBD followed by infection with the SARS-CoV-2 at an MOI of 0.5. The cells were stained for spike protein and the percentage of cells expressing the spike protein in each condition was plotted. Representative data of CBD from Figure 1C (Supplier A) is used for comparison. EC50 values are indicated.

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 2
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 3. CBD inhibits viral replication after SARS-CoV-2 entry into the host cell. (A) Immunoblots of ACE2 protein expression from A549-ACE2 cell lysates either untreated or treated with vehicle or CBD at indicated doses (n = 3). Blots were probed with antibodies against ACE2 and tubulin. ACE2 protein expression levels were normalized to the tubulin signal within each sample. ACE2 expression levels were plotted relative to untreated samples. (B) 293 T-ACE2 cells were infected by spike or VSV-G pseudovirus for 72 hours with the indicated doses of CBD treatment, and the percentage of infected cells plotted. (C) A549-ACE2 cells were infected with SARS-CoV-2 at an MOI of 0.5 for 2 hours. DMSO or 10 μM CBD was then added at either 2, 6 or 15 hours after infection. After 16 hours, spike positive cells were quantified and normalized to the virus-infected only samples. (D) Left panel: A549-ACE2 cells were infected with SARS-CoV-2 at an MOI of 0.5 for 2 hours. DMSO or 10 μM CBD was then added at 2 hours after infection with the spike neutralizing antibody to prevent reinfection. After 16 hours, spike positive cells were quantified and normalized to the virus-infected only samples. Right panel: Validation of neutralizing antibody efficacy. 400 pfu of SARS-CoV-2 virus was incubated with or without 100 μM of neutralizing antibody for 1 hour. A549-ACE2 cells were treated with the mixture for 16 hours and Spike positive cells were quantified.

CBD inhibits viral RNA expression and reverses viral-induced changes in host gene expression

F4 Consistent with this interpretation, RNA-seq analysis of infected A549-ACE2 cells treated with CBD for 24 hours shows a striking suppression of SARS-CoV-2-induced changes in gene expression. CBD effectively eradicated viral RNA expression in the host cells, including RNA coding for spike, membrane, envelope and nucleocapsid proteins (Figs. 4 A and B). Both SARS-CoV-2 and CBD each induced significant changes in cellular gene expression (figs. S5 and S6). Principal component analysis (PCA) of host cell RNA shows almost complete reversal of viral changes but, rather than returning to a normal cell state, the CBD + virus infected cells resemble those treated with CBD alone (Fig. 4C). Clustering analysis using Metascape reveals some interesting patterns and associated themes (Fig. 4D, figs. S7, and S8). For example, viral induction of genes associated with chromatin modification and transcription (Cluster 1) is reversed by CBD, although CBD alone has no effect. Similarly, viral inhibition of genes associated with ribosomes and neutrophils (Cluster 3) is largely reversed by CBD, but the drug alone has no effect. This contrasts with Clusters 5 and 6 where CBD alone induces strong activation of genes associated with the host stress response. Together

these results suggest that CBD acts to prevent viral protein translation and associated cellular changes.

To gain a better understanding of the specific anti-viral action of CBD, we analyzed RNAseq from lysates of uninfected or SARS-CoV-2-infected cells treated for 24 hours with the inactive CBDV homologue. Induction of viral genes for spike, envelope and nucleocapsid proteins is reduced by only 60% with CBDV as opposed to ~99% with CBD (Fig. 5A,B). CBDV treatment causes fewer transcriptomic changes than CBD in A549-ACE2 cells and is largely ineffective at reversing transcriptional changes induced by SARS-CoV-2 (Fig. 5C). Clustering analysis using Metascape reveals only a couple clusters that show CBDV reversal of viral transcriptomic changes (Fig. 5D). These include autophagy and lipid metabolism (Cluster 1) that are induced by CBDV as well as protein translation/cell cycle/DNA replication (Cluster 3) that are suppressed by CBDV.

CBD induces the ER stress response and IRE1α activity as a key mechanism for its anti-viral action

Of particular interest are three sets of genes related to the endoplasmic reticulum (ER) stress response, the unfolded protein response (UPR) and interferon induction that are selectively upregulated by CBD

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 3
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 4. Changes in Viral and host cell transcription following SARS-CoV-2 infection or CBD treatment. A549-ACE2 cells were infected with SARS-CoV-2 at MOI of 3 with or without CBD treatment at 10 μ M for 24 hours. RNA-seq was performed as described in Methods. **(A)** Heatmap of relative levels of SARS-CoV-2 genes from the RNA-seq samples. **(B)** Expression levels of SARS-CoV-2 spike and nucleocapsid genes. Percent expression level changes for genes from infected cells compared to cells infected and CBD treated are indicated for each gene. **(C)** Principal component analysis (PCA) of RNA-seq data showing control (veh_mock), SARS-CoV-2 infected (veh_infect), CBD-treated (CBD_mock), and SARS-CoV-2 infected plus CBD treated (CBD_infect) samples. The first and second principal components (PC1 and PC2) of each sample are plotted. **(D)** Heatmap of normalized expression levels of 5,000 most variable genes across all RNA-seq samples, clustered into 6 groups based on differential expression between treatment conditions.

F6 but not CBDV (Fig. 6A). By contrast, genes associated with the oxidative stress response are induced by both cannabinoids. Cells experience ER stress when the workload on the ER protein folding machinery exceeds its capability. Under ER stress, secretory proteins accumulate in unfolded forms within the organelle to trigger a set of intracellular signaling pathways called UPR, which is part of a larger cellular stress response that maintains proteostasis throughout the cell (18). The UPR pathway is controlled by three ER transmembrane proteins – IRE1 α , PERK, and ATF6 – that contain an ER luminal domain capable of directly or indirectly sensing misfolded proteins. In response to ER stress, each of these sensors sets in motion transcriptional and translational changes that increase protein folding capacity and attempt to restore homeostasis. However, if the stress on the ER is irremediable, the UPR switches outputs and signals cell death. We validated CBD induction of IRE1 α , PERK, and ATF6 gene expression by qRT-PCR (fig. S9A), consistent with previous reports (19). Ingenuity analysis confirmed that CBD induces the UPR significantly more than CBDV (figs. S9B, S10B, S11).

Numerous studies report compelling evidence that the UPR is hyperactivated and required for replication of other closely related

coronavirus family members (20, 21). Surprisingly, although GSEA enrichment analysis of the RNA-seq data showed that the IRE1 α pathway is strongly activated by CBD in the presence or absence of virus, this pathway was not activated by SARS-CoV-2 alone (Table 1; figs. S12, S13, S14). PERK, by contrast, was functionally activated by both SARS-CoV-2 and CBD. IRE1 α is a single pass ER transmembrane protein with bifunctional kinase/endoribonuclease (RNase) activities. In response to ER stress, IRE1 α undergoes oligomerization and autophosphorylation, which allosterically activates its RNase to initiate productive splicing of XBP1 mRNA. Spliced XBP1 encodes a transcription factor that upregulates many host stress responses, including ER chaperone induction and ER-associated degradation (ERAD) components (22) (Fig. 6E).

CBD strongly activates IRE1 α RNase activity as shown by analysis of XBP1 splicing using both RNaseq data to quantify spliced XBP1 as well as direct confirmation by qRT-PCR (Fig. 6B; fig. S15). As predicted, CBD induced XBP1 splicing in the presence or absence of virus whereas CBDV had no significant effect and is comparable to virus alone. The time course and dose response curves for CBD induction of XBP1 splicing in the absence of the virus were

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 4
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 5. Changes in Viral and host cell transcription following SARS-CoV-2 infection or CBDV treatment. A549-ACE2 cells were infected with SARS-CoV-2 at MOI of 3 with or without CBDV treatment at 10 μ M for 24 hours. RNA-seq was performed as described in Methods. **(A)** Heatmap of relative levels of SARS-CoV-2 genes from the RNA-seq samples. **(B)** Expression levels of SARS-CoV-2 spike and nucleocapsid genes. Percent expression level changes for genes from infected cells compared to cells infected and CBD treated are indicated for each gene. **(C)** Principal component analysis (PCA) of RNA-seq data showing control (veh_mock), SARS-CoV-2 infected (veh_infect), CBDV-treated (CBDV_mock), and SARS-CoV-2 infected plus CBDV treated (CBDV_infect) samples. The first and second principal components (PC1 and PC2) of each sample are plotted. **(D)** Heatmap of normalized expression levels of 5,000 most variable genes across all RNA-seq samples, clustered into 6 groups based on differential expression between treatment conditions.

consistent with the time course and dose responses for CBD inhibition of viral spike protein expression in A549-ACE2 cells (Fig. 6C). Furthermore, while an IRE1 α knockout had no significant effect on SARS-CoV-2 infection, it shifted the dose response and significantly reduced the anti-viral effects of CBD, leading to an approximately 2-fold increase in its EC50 against SARS-CoV-2 (Fig. 6D; fig. S16). Together, these results indicate that CBD induction of IRE1 α is a critical component of its anti-viral action against SARS-CoV-2.

CBD induces interferon expression as part of its anti-viral activity

Another mechanism by which CBD could suppress viral infection and promote degradation of viral RNA is through induction of the interferon signaling pathway. Interferons are among the earliest innate immune host responses to pathogen exposure (23). As reported (24), SARS-CoV-2 infection suppresses the interferon signaling pathway (Fig. 7A, and fig. S17). Many genes in the pathway such as ISG15, IFIT1, IFIT3, SOCS1 and OAS1, an interferon-induced gene that leads to activation of RNase L and RNA degradation (25),

were moderately up-regulated by CBD alone but highly induced by CBD in the presence of the virus (Fig. 7A and figs. S18,19). These latter results are consistent with the possibility that CBD lowers the effective viral titer sufficiently to enable normal host activation of the interferon pathway. At the same time, CBD effectively reversed viral induction of cytokines that can lead to the deadly cytokine storm at later stages of infection (Fig. 7B). By contrast, the inactive homologue CBDV does not significantly induce genes within the interferon pathway or prevent cytokine induction (Fig. 6A, 7A, 7C, figs S20A,B and S21).

To directly test the possibility that interferons might account in part for the anti-viral activity of CBD, we exposed ACE2-A549 cells to a mixture of antibodies against Type I (α,β,ω) and Type II (γ) interferons prior to 2.5 μ M CBD treatment and viral infection. The results show that the anti-interferon antibodies reduce the anti-viral effects of CBD and partially rescue SARS-CoV-2 infection (Fig. 7D). Collectively, these results suggest that CBD inhibits SARS-CoV-2 infection in part by activating IRE1 α and the interferon pathways, leading to degradation of viral RNA and subsequent viral-induced changes in host gene expression, including cytokines.

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 5
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 6. CBD promotes host cell ER stress responses and IRE1 α /XBP1 splicing, and IRE1 α contributes to anti-viral CBD activity. A549-ACE2 cells were infected with SARS-CoV-2 at MOI of 3 with or without CBD or CBDV treatment at 10 μ M (A,B) or as indicated (C) for 24 hours (A, B) or as indicated (C). (A) Heatmap of predicted pathway activation based on Ingenuity analysis of activation z-scores for each pathway and each comparison. Red: pathway is activated. Blue: pathway is inhibited. White: pathway is unchanged. Gray: no prediction due to lack of significance. (B) Analysis of XBP1 splicing by the IRE1 α RNase. Reads representing spliced or unspliced XBP1 were identified and quantified for cells that were mock-treated, SARS-CoV-2 treated or treated with CBD or CBDV either alone or in the presence of virus (left panel). Percentage of alternatively spliced reads for the RNA-seq samples were plotted and unpaired t tests were performed comparing each experiment's mock samples to other samples (right panel). (C) A549-ACE2 cells were treated by indicated concentrations of CBD for 3 and 6 hours. Relationship between CBD concentration and XBP1 splicing were determined by qRT-PCR. (D) Effect of IRE1 α on dose response for anti-viral activity of CBD. A549-ACE2 or A549-ACE2 cells lacking IRE1 α (IRE1 KO) were treated with indicated doses of CBD followed by infection with SARS-CoV-2 at an MOI of 0.5 for 48 hours. The cells were stained for spike protein and the percentage of cells expressing the spike protein in each condition was plotted. EC50 values are indicated. Unpaired t tests were performed at each concentration and significant p-values were shown. This is representative of three independent experiments (Composite EC50 1.7 v. 1.2, $p < 0.05$). (E) Schematic illustrating effect of CBD and SARS-CoV-2 on IRE1 α RNase activity and XBP1 splicing.

CBD treatment significantly inhibits SARS-CoV-2 replication in mice

As several agents including cationic amphipathic drugs block SARS-CoV-2 replication in cultured cells but not *in vivo* (26), we determined whether CBD reduces viral titer in female K18-hACE2 mice (27). Mice were injected intraperitoneally twice daily with CBD (20 or 80 mg/kg) for 7 days prior to intranasal challenge with SARS-CoV-2 (2×10^4 PFU). After the challenge, administration of CBD continued twice daily for an additional 4 days (Fig. 8A). CBD treatment significantly inhibited viral replication in lungs and nasal turbinates at day 5 post-infection in a dose-dependent manner (Figs. 8B-C). The lower dose of CBD reduced viral load by 4.8-fold

in lungs and 3.7-fold in nasal turbinates, while the higher dose decreased viral titers by 40- and 4.8-fold in lungs and nasal turbinates, respectively. During this period, the mice showed no signs of clinical disease, and their body weights were not significantly changed (Fig. 8D). These results establish the pre-clinical efficacy of CBD as an anti-viral drug for SARS-CoV-2 during early stages of infection.

CBD usage is negatively associated with indications of SARS-CoV-2 infection in patients

Given that high purity CBD preparations are taken by a large number of individuals, we examined whether medication records of CBD prescriptions or use are associated with indications of SARS-CoV-2

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 6
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

Table 1. Induction of PERK, IRE1 or ATF6 gene expression and function in response to CBD and/or SARS-CoV-2 virus. RNA-seq gene expression data are used for GSEA on three GO terms “PERK-mediated UPR”, “IRE1-mediated UPR” and “ATF6-mediated UPR” (Gene Ontology numbers 36498, 36499, 36500). Normalized enrichment score is shown under the “GSEA NES” column (higher score = more enrichment). Fold-change for transcriptional expression differences between PERK, IRE1, and ATF6 is shown for each comparison under the “RNA-seq fold-change” column.

Comparison	UPR branch	GSEA NES	RNA-seq fold-change
CBD vs mock	PERK	1.43	2.46
	IRE1	1.38	2.29
	ATF6	ND*	1.40
Virus vs mock	PERK	1.92	1.85
	IRE1	Not enriched	2.67
	ATF6	ND*	0.91
CBD + virus vs mock	PERK	1.45	3.24
	IRE1	1.44	2.76
	ATF6	ND*	1.24

*ND = Not determined due to not enough genes to get reliable values.

infection (i.e., positive COVID-19 tests and/or COVID-19 diagnoses proximal to COVID-19 tests). An oral solution of CBD 100 mg/mL (CBD100) is often used for the treatment of seizures (see the Patient Analysis Supplement). Analysis of 1,212 patients from the National COVID Cohort Collaborative (N3C) (28) with a history of seizure-related conditions and a medication record of CBD100 revealed 6.2% (75 patients) with an indication of SARS-CoV-2 infection proximal to the dates of their first COVID-19 test in their N3C data. This was a significantly lower rate than the rates of matched control groups of patients that did not have any CBD100 records (e.g., 6.2% for CBD100 patients compared to 8.9% for non-CBD100 patients, $p = 0.014$; multivariable logit model odds ratio (OR) of 0.65, $p = 0.009$, 95% C.I. [0.47,0.90]). The demographics and medication history of the CBD100 patients were similar to those of the matched control group. The medical condition history for these patients included seizure-related conditions, the CDC list of at-risk conditions (29) and other potential confounders such as conditions of reduced mobility, chronic pain, or developmental disabilities that can limit public interaction and COVID-19 exposure. The negative association was even more significant in analyses of a subgroup of 531 CBD100 patients who were likely taking CBD100 on the dates of their first COVID-19 tests (e.g., 4.9% among these CBD100 patients compared to 9.0% among 531 matched controls, $p = 0.011$; OR = 0.48, $p = 0.006$, 95% C.I. [0.29,0.81]) (Fig. 9; Table S4 in Patient Analysis Supplement which describes the patient data analysis methods and findings in detail).

DISCUSSION

Our results suggest that CBD and its metabolite 7-OH-CBD can block SARS-CoV-2 infection at early and even later stages of infection. The mechanism appears to be mediated in part by activation of the IRE1 α RNase and interferon pathways. In addition to these

cell-based findings, pre-clinical studies show that CBD treatment reduced viral titers in the lungs and nasal turbinates of SARS-CoV-2-infected mice. Finally, analysis of a national sample of patients with active records of 100 mg/ml CBD consumption at the time of COVID testing revealed an association with substantially fewer SARS-CoV-2 positive test results. This negative association was robust to many sensitivity analyses, including changes in the matching and outcomes models, and merits further research into the potential of CBD to combat SARS-CoV-2 infection, such as validation in other large, multi-site electronic health record datasets or prospective experimental designs.

One mechanism contributing to the antiviral activity of CBD is the induction of the interferon pathway both directly and indirectly following activation of the host immune response to the viral pathogen. In fact, interferons have been tested clinically as potential treatments for COVID-19 (30). When hyperactivated by severe ER stress, IRE1 α 's RNase activity leads to the endonucleolytic decay of many ER-localized mRNAs (RIDD) and subsequent activation of RIG-I and interferons (18). Although SARS-CoV-2 induces the kinase activity of IRE1 α , it does not activate its RNase activity as monitored by XBP1 splicing. Thus, the RNase activity of IRE1 α induced by CBD can potentially account both for the degradation of viral RNA and the induction of interferons by the RNA fragments. Further investigation will be required to determine whether both anti-viral effects of CBD are linked to the ER stress response. Importantly, CBD also suppresses cytokine activation in response to viral infection, reducing the likelihood of immune cell recruitment and subsequent cytokine storms within the lungs and other affected tissues. These results complement previous findings suggesting that CBD suppresses cytokine production in recruited immune cells such as macrophages (31). Thus, CBD has the potential not only to act as an antiviral agent at early stages of infection, but also to protect the host against an overactive immune system at later stages.

CBD has a number of advantages as a potential preventative agent against SARS-CoV-2. CBD as a food additive with THC content less than 0.3% is widely available without restricted access. With proper formulation, quality control and delivery, CBD could be used prophylactically in contrast to recent anti-viral drugs. Multiple means of CBD ingestion are possible, including potential for inhalation and nasal delivery. CBD blocks viral replication after entry into cells and, thus, is likely to be effective against viral variants with mutant spike proteins. Unlike drugs such as remdesivir or antiviral antibodies, CBD administration does not require injection in hospital settings. Finally, CBD is associated with only minor side effects (32).

However, several issues require close examination before CBD can be considered further or even explored as a therapeutic lead for COVID-19 (12). Although many CBD and CBD-containing products are available on the market, they vary vastly in quality, CBD content, and their pharmacokinetic properties after oral administration, which are mostly unknown. CBD is quite hydrophobic and forms large micellar structures that are trapped and broken down in the liver, thereby limiting the amount of drug available to other tissues after oral administration. Inactive carriers and formulation adjuvants have a significant impact on clinically obtainable concentrations. As CBD is widely sold as a preparation in an edible oil, we analyzed flavored commercial hemp oils and found a CBD content of only 0.30% in a representative sample (fig. S22). The purity of CBD and the chemical composition of the materials labelled as CBD are also important, especially in light of our findings suggesting that other cannabinoids

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no:
---	--------------------------	-----------------------------------	-------	-------	-----------	---------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 7. CBD promotes host cell interferon responses and inhibits viral induction of cytokines. (A) Heatmap of fold-change (log₂) of genes from the Interferon Response Canonical Pathway for all virus or CBD-treated samples compared to mock-treated samples. Columns 1–3 use samples from the RNA-seq experiment on CBD and SARS-CoV-2. Columns 4–6 use samples from the RNA-seq experiment on CBDV and SARS-CoV-2. (B) Heatmap of normalized expression levels of GO Cytokine Activity genes which were up-regulated by the viral infection but down-regulated by CBD treatment for all RNA-seq samples from the experiment on CBD and SARS-CoV-2. (C) Heatmap of the normalized expression levels of the same genes for all RNA-seq samples from the experiment on CBDV and SARS-CoV-2. (D) A549-ACE2 cells were treated with 2.5 μM of vehicle or CBD with or without Human IFN-γ Antibody and Human Type I IFN Neutralizing Ab Mixture at 2 hours before infection. Cells were then infected with 0.5 MOI SARS-CoV-2 and incubated for 24 hours, and active virus was measured using a plaque assay. The results are representative of three independent experiments.

such as THC might act to counter CBD antiviral efficacy. This essentially eliminates the feasibility of marijuana serving as an effective source of antiviral CBD, in addition to issues related to its legal status. Finally, other means of CBD administration such as vaping and smoking raise additional concerns about potential lung damage.

Future studies to explore the optimal means of CBD delivery to patients along with clinical trials will be needed to further evaluate the promise of CBD as a therapeutic to block SARS-CoV-2 infection. Our animal studies provide pre-clinical support for evaluation of CBD as an anti-SARS-CoV-2 therapeutic agent in clinical trials. We

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 7
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 8. CBD inhibits SARS-CoV-2 replication in mice. (A) Timeline of the mouse experiment. (B) Viral titer in lungs from all animals measured 5 days after viral challenge (day 12). (C) Viral titer in nasal turbinates from all animals measured 5 days after viral challenge (day 12). (D) Weight measurements of mice in each treatment group (n = 10) during the study. The body weight of each mouse is normalized to its weight measured at Day 0. All animals were challenged with SARS-CoV-2 by intranasal instillation at Day 7.

advocate carefully designed placebo-controlled clinical trials with known concentrations and highly-characterized formulations in order to define CBD's role in preventing and treating early SARS-CoV-2 infection. The necessary human *in vivo* concentration and optimal route and formulation remain to be defined. We strongly caution against the temptation to take CBD in presently available formulations including edibles, inhalants or topicals as a preventative or treatment therapy at this time, especially without the knowledge of a rigorous randomized clinical trial with this natural product (33).

MATERIALS AND METHODS

Study design

The goal of this study was to determine whether cannabidiol (CBD), a natural product extracted from the cannabis plant, has the potential to inhibit infection of cells by SARS-CoV-2. To this end, we utilized three different human or monkey cell lines. We tested four independent preparations of CBD from chemical as well as natural sources, and also tested related cannabinoid compounds and metabolites. We used RNA-seq analysis to demonstrate that CBD, in contrast to the inactive cannabinoid CBDV, effectively eliminated SARS-CoV-2 viral RNA from infected cells, activated the ER stress response and XBP1 splicing, induced expression of the interferon pathway and suppressed viral induction of cytokines. We demonstrated

using IRE1 α knockout cells and anti-interferon blocking antibodies that both IRE1 and interferons contribute to the anti-viral activity of CBD. Finally, utilizing medical records for groups of human patients from the National COVID Cohort Collaborative under appropriate IRB protocols, we analyzed the association of patients taking CBD with their risk of testing positive for SARS-CoV-2. Statistics are provided in the corresponding figures and in methods.

Materials, cells and viruses

High-purity CBD was acquired from two chemical companies or two online commercial sources. 7-OH-CBD was purchased from Cerilliant Corporation (Round Rock, TX). All commercial compounds used were validated by NMR as described below. Cannabinoid-infused hemp oil containing 1,500+ mg cannabinoids was from Bluebird Botanicals (Louisville, CO, USA). Hemp extract from *C. sativa* biomass was from Hopsteiner Ltd. (Yakima, Washington, USA). Low CBD hemp oil was obtained from an online commercial source. A549-ACE2 cells were generously provided by tenOever and colleagues (24). Vero E6 cells and Calu3 cells were purchased from ATCC. SARS-CoV-2 (nCoV/Washington/1/2020) was provided by Natalia Thornburg (Centers for Disease Control) via the World Reference Center for Emerging Viruses and Arboviruses (Galveston, Texas), and from BEI Resources for the *in vivo* studies. SARS-CoV-2 variants were provided by BEI resources. The α variant is BEI number

MS no:	Emp name:	Date / Time:	PE's:	AA's:	Comments:	Art no:
RAabi6110/IMMUNOLOGY, LIFE SCIENCES	jrnindesign	1-20-2022 / 22:52				Fig 8

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.




Fig. 9. CBD 100 mg/mL medication records in patients are significantly associated with less COVID-19 positivity. Schematic showing derivation of our Main Analysis Sample and cannabidiol (CBD) patient groups obtained from the National COVID Cohort Collaborative (N3C). Successive analyses of patient subsets are illustrated. The final panel shows associations between having a CBD 100 mg/mL medication record on the date of their first COVID-19 test and COVID-19 Positive Status among matched control groups of increasing size (i.e., 1-to-1, 2-to-1, and 3-to-1 ratios of controls to CBD patients). A Mismatched Covariate has a standardized mean difference greater than 0.10 and a two-sided Fisher exact test p-value less than 0.05 when comparing its distribution between the CBD patients and their matched controls. AUC refers to area under the receiver operating characteristic curve. Detailed information regarding the patient data analysis methods and findings is in the Patient Analysis Supplement.

NR-54000, isolate hCoV-19/England/204820464/2020 sourced from Public Health England. The β variant is BEI number 54009, B.1.351(20H/501Y.V2) sourced from the Africa Health Research Institute. The γ variant is BEI number 54982, isolate hCoV-19/Japan/TY7-503/2021 sourced from the Japan National Institute of Infectious Disease. Viral stocks were made by two passages in Vero E6 cells and stock titers were determined by limiting dilution plaque titer on VeroE6 cells (described below).

SARS-CoV-2 infection assay

All SARS-CoV-2 infections were performed in biosafety level 3 conditions at the Howard T. Rickett Regional Biocontainment Laboratory, University of Chicago. *In vivo* infections were performed in ABSL-3 conditions at the Center for Predictive Medicine for Biodefense and Emerging Infectious Diseases, the University of Louisville Regional Biocontainment Laboratory. Cells in DMEM +2% FBS were treated with CBD or other inhibitors or 2 hours with

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no: Fig 9
---	--------------------------	-----------------------------------	-------	-------	-----------	------------------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

2-fold dilutions beginning at 10 μM in triplicate for each assay. A549-ACE2 cells were infected with an MOI (multiplicity of infection) of 0.5 in media containing the appropriate concentration of drugs. Vero E6 cells were infected with an MOI of 0.1 in media containing the appropriate concentration of drugs. After 48 hours, the cells were fixed with 3.7% formalin, blocked, and probed with mouse anti-Spike antibody (GTX632604, GeneTex) diluted 1:1,000 for 4 hours, rinsed, and probed with anti-mouse-HRP for 1 hour, washed, then developed with DAB substrate 10 minutes. Spike positive cells ($n > 40$) were quantified by light microscopy as blinded samples. Viral titers were determined by plaque assay. Briefly, a monolayer of E6 cells is infected with a series of serial dilutions of virus sample for 1 hour at 37°C. The viral inoculum is then removed and replaced by a MEM overlay media containing 1.25% carboxymethyl cellulose. Cells are incubated for 72 hours after which overlay media is removed and cells are fixed with 10% formalin and stained with 0.25% crystal violet solution. Plaques are counted in the dilution well with between 10–100 plaques and original concentration of viral sample is calculated. Data were analyzed and plotted using GraphPad Prism and EC50 values were extracted from nonlinear fit of response curves.

Crystal Violet toxicity assay

Cells were treated with varying concentrations of different compounds in 2% DMEM starting at 10 μM and going down by 1/2 for six more dilutions. Cells were incubated with the drug for 48 hours. Cells were fixed with 10% Formalin solution for 30 minutes. Then they were stained with 1% Crystal Violet solution for 30 minutes after which plates were dried and the amount of Crystal Violet staining was assessed by measuring absorbance at 595 nm on a TECAN M200 Plate reader. Absorbance readings were normalized to those of the control wells not treated by the drug to measure the differences in cell growth with or without the drug treatment.

Spike protein and antibody neutralizing assay

A549-ACE2 cells were treated with 10 μM of CBD either 2 hours before infection or 2, 6 or 15 hours after infection. Cells were infected with MOI of 0.5 for 2 hours. Then, the infection media was replaced with media containing CBD or DMSO, and the samples were incubated at 37°C for 16 hours. In one experiment when CBD was added 2 hours after infection, infection media was replaced with CBD or DMSO and μM of neutralizing antibody (Active Motif 001414). After 16 hours, the samples were fixed with 10% formalin and underwent IHC for spike protein. Neutralizing antibody efficiency was tested by incubating 400 pfu of virus with or without 100 μM of the antibody at 37°C for 1 hour. Then A549-ACE2 cells were infected with the mixture for 16 hours. Spike positive cells were quantified as described above.

Interferon antibody neutralizing assay

A549-ACE2 cells were treated with 2.5 μM of CBD, 1 $\mu\text{g}/\text{ml}$ Human IFN- γ Antibody (MAB285-100) and 1:25 dilution of Human Type I IFN Neutralizing Ab Mixture (PBL Assay Science 39000-1) 2 hours before infection. Cells were then infected with 0.5 MOI and incubated for 24 hours, after which supernatants were collected and active virus measured using plaque assay described above.

Generation of IRE1 α knockout cells by CRISPR-Cas9

Lentivirus stocks were by using lentiCRISPR v2 (Addgene) with single guide RNA (sgRNA) targeting IRE1 sequence (CGGTCACT-CACCCCGAGGCC). The infected A549-ACE2 cells were polyclonally

selected and maintained by using medium supplemented with 4 $\mu\text{g}/\text{mL}$ puromycin for 1 week.

Description of the cannabinoids

CBD can be procured by isolating cannabidiolic acid (CBDA) from *Cannabis sativa* plant material and then inducing chemical decarboxylation, or via decarboxylation of cannabinoids contained in raw plant material or extract and subsequent isolation of CBD. Cannabidiol (CBD) is a naturally occurring CBD homolog that has an *n*-propyl in place of CBD's *n*-pentyl side chain. Cannabigerol (CBG), in the form of cannabigerolic acid, is the metabolic precursor to both tetrahydrocannabidiolic acid and CBDA in *C. sativa*. Tetrahydrocannabinol (THC) is a cyclized congener of CBD that is obtained after tetrahydrocannabinolic acid decarboxylation. THC is present in *C. sativa* in both Δ^9 -*cis* and Δ^9 -*trans* stereoisomers. Cannabichromene (CBC), in the form of cannabichromenic acid, represents a third possible cannabigerolic acid metabolite with a chromene ring in the geranyl residue.

Acquisition, isolation and characterization of cannabinoids

In the present study, purification of CBD from natural sources used (a) cannabinoid infused hemp oil containing 1,500+ mg cannabinoids in medium-chain triglycerides per fluid ounce, manufactured by Bluebird Botanicals (Louisville, CO, USA), and (b) hemp extract prepared by supercritical fluid extraction (SFE) with CO₂ from *C. sativa* biomass qualifying as hemp, manufactured by Hopsteiner Ltd. (Yakima, Washington, USA) with a 54.7% total content of CBD, calculated as CBD + CBDA*0.877. Typical purities of these CBD preparations are in the 90–97% range including foreign impurities (e.g., residual solvent) determined by qHNMR. Details of the purification and structure analysis methodologies are detailed in a concurrent publication, which is currently in press (Journal of Natural Products). In brief, the methodologies can be summarized as follows:

Purification Procedure

CBD, CBC, CBG, Δ^9 -*trans*-THC, Δ^9 -*cis*-THC, and CBDV were isolated from the hemp oil, and CBDA from the crude hemp SFE extract, using centrifugal partition chromatography (CPC), a countercurrent separation technique, and a biphasic liquid-liquid solvent system.

Structure Elucidation Methodology

The identities of the commercially sourced CBD and other cannabinoid samples were verified by 1D ¹H NMR analysis, performed as qNMR measurement, via comparison with an authentic HfSA profile of CBD as published (12). In addition to an overall excellent match of the profiles, the highly coupled fingerprint signal of H-4"ax served as a highly specific identity marker. The structures of the cannabinoids that were sourced commercially or purified from the natural sources were established by a combination of 1D/2D NMR and LC-HRMS analysis, taking into account reference data from the literature.

NMR Sample Preparation

For commercial samples supplied as solution, the solvent was removed carefully in vacuo and 450 μL deuterated methanol (MeOH-*d*₄) added to the residue using a precision syringe. The solution was transferred into 5-mm NMR tube with a glass pipette, the vial rinsed three times with 25 μL of solvent and the rinsing solution transferred into the same NMR tube, for a final volume of 525 μL . Commercial and isolated samples available as solids were directly weighed into a 5-mm NMR tube and 500 μL of solvent added with a precision syringe. For analysis of the commercial hemp oil preparation, 10 drops (0.25 mL equivalent to 14–15 drops) was added into 5 mm

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no:
---	--------------------------	-----------------------------------	-------	-------	-----------	---------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

NMR tube directly. The net weight of hemp oil in NMR tube was 198.50 mg, determined on a 0.01 mg precision balance, and 0.90 mg of dinitrobenzoic acid were added as an internal calibrant for IC-qHNMR; 325 μ L of CDCl_3 and 10 μ L of CD_3OD were added, and the tube was flame sealed.

NMR data acquisition and processing and qNMR evaluation

All ^1H NMR data were acquired on a Bruker 600 Avance III with a two channel ^{13}C direct cryogenic probe. Time domain (TD) was set to 64 k, relaxation delay (D1) was 60 sec, 90-degree excitation pulses were used for a total of 32 signal averaged scans. The receiver gain (RG) was 32 for all samples, except for one mass-limited sample < 1 mg (RG = 101) and the large-quantity hemp oil sample (RG = 2; 15 degrees excitation pulse used). Determination of sample purity and CBD content in hemp oil by quantitative NMR (qNMR) using the 100% qNMR approach and openly published worksheets (<https://gfp.people.uic.edu/qnmr/content/qnmrcalculations/100p.html>). The qNMR purity of all CBD samples was >97% including foreign impurities, and no cannabinoid congeners could be detected at levels above 1.0%. Using the absolute qHNMR method with internal calibration (IC abs-qNMR), the content of CBD in hemp oil was determined as 0.30%.

Pseudotyped lentivirus production

293 T and 293 T-ACE2 cells were cultured in DMEM (Corning 10017CV) with 1X sodium pyruvate (Gibco 11360070) and 10% FBS (HyClone SH30910). Lentivirus particles pseudotyped with SARS-CoV-2 (Wuhan-Hu-1) spike protein or VSV-G, were generated as described (19). Briefly, 293 T cells were transfected using TransIT-LT1 (Mirus) with third generation lentivirus packaging vectors (HDM-Hgpm2, HDM-tat1b, pRC-CMV-Rev1b), transfer vector (pHAGE-CMV-ZsGreen-W) and either SARS-CoV-2 spike (HDM-IDTSpike-fixK) or VSV-G (HDH-VSVG). Supernatants collected at 36 and 60 hours post-transfection were pooled, syringe filtered and frozen in single-use aliquots at -80°C . All plasmids used for lentivirus production were kindly provided by Dr. Jesse Bloom (University of Washington, Seattle).

Pseudovirus binding assay

293 T-ACE2 cells were seeded at 1.2×10^4 cells per 96well in black wall, clear bottom plates. The next day, 2-fold dilutions of CBD stock (10 mM) were prepared in DMSO, followed by 1:1000 dilutions in either complete DMEM or pseudovirus preparation. SARS-CoV-2 spike pseudovirus was used undiluted, while VSV-G pseudovirus was diluted 1:1,500 in complete DMEM. Cells and pseudovirus were pre-treated with CBD dilutions for 2 hours and 1 hour at 37°C , respectively. Cells were infected with pseudovirus for 72 hours, fixed with 4% paraformaldehyde, stained with a nuclear marker (Hoechst 33342, ThermoFisher H3570) and imaged. 293 T-ACE2 cells were generously supplied by Dr. Jesse Bloom (University of Washington, Seattle).

Pseudovirus neutralization assay

293 T or 293 T-ACE2 cells were seeded at 1.2×10^4 cells per 96-well in black wall, clear bottom plates. Next day, SARS-CoV-2 spike neutralizing antibody (Sino Biological 40592-R001) was diluted in complete DMEM to a starting final concentration of 300 ng per 100ul per 96well, followed by subsequent 3-fold dilutions. Neutralizing antibody was incubated with pseudovirus for 1 hr. at 37°C . Cells

were infected with pseudovirus +/- neutralizing antibody for 72 hrs, fixed with 4% paraformaldehyde, stained with nuclear marker Hoechst 33342 and imaged.

Protease inhibition assay

Assays were performed in duplicate at room temperature in 96-well black plates at 25°C . Reactions containing varying concentrations of inhibitor (10 or 50 μM) and 3CLpro enzyme (0.4 μM) or PLpro enzyme (0.3 μM) in Tris-HCl pH 7.3, 1 mM EDTA were incubated for approximately five minutes. 3CLpro reactions were then initiated with TVLQ-AMC probe substrate (40 μM) and PLpro reactions were initiated with LKGG-AMC probe substrate (40 μM). The reaction plate was shaken linearly for 5 s and then measured for fluorescence emission intensity (excitation λ : 364 nm; emission λ : 440 nm) over time (1 min-3 h) on a Synergy Neo2 Hybrid. Each assay contained 2-3 positive control wells (DMSO) and 2 negative control wells (assay components without protease). Data were normalized to the positive control wells at 3 h, which was assigned an arbitrary value of 100.

Immunoblotting

A549-ACE2 cells were treated with CBD, vehicle (DMSO) or not treated for 24 hours. Cells were first washed with ice-cold PBS. Whole-cell extraction were prepared by directly lysing cells with Laemmli sample buffer (Bio-rad 1610747) supplemented with protease inhibitor (Roche 4693159001), PMSF (Roche 10837091001) and phosphatase inhibitor (GB-450) at 4°C . Protein samples were finally boiled at 98°C for 5 mins. Western blotting was performed using antibodies for ACE2 (Abcam 108252) and α -tubulin (Invitrogen MA1-19401) for control. For validations of IRE1 α knockout in A549-ACE2, cells, antibodies for IRE1 α (Cell Signaling, 3294S) and GAPDH (Cell Signaling, 5174S) were used. Blots were imaged and quantified using Licor Odyssey Fc.

RNA sequencing

Lung alveolar A549 cells were stably overexpressed with human angiotensin converting enzyme 2 (ACE2) protein and seeded at 10,000 cells per well in a 96-well plate. Cannabidiol or vehicle were added together to the cells. Cannabidiol (Cayman Chemical, 90080) was dissolved in a 10 mM stock solution with DMSO (Sigma-Aldrich, D2650-100 mL). Final concentration of CBD was 10 μM . The virus stock was then removed and replaced with fresh 2% FBS DMEM media with drug. The cells were incubated for another 24 hours before total RNA extraction the NucleoSpin 96 RNA kit (Takarabio, 740709). Three independent biological replicates were performed per experimental condition, with 12 total RNA samples. RNA sample quality check, library construction, and sequencing were performed by the University of Chicago Genomics Facility following standard protocols. The average RNA Integrity Score was 8.9. All 12 samples were sequenced in two runs by a NovaSeq 6000 sequencer to generate paired-end 100 bp reads. For each sample, the raw FASTQ files from two flow cells were combined before downstream processing. Cannabidiol (CBDV) was isolated from the hemp oil as described above, and identical studies as those described above with CBD were performed. The average RNA Integrity Score for the CBDV samples was also 8.9.

RNA-seq data for both CBD and CBDV treated cells were analyzed separately using a local Galaxy 20.05 instance for the following steps (34). Quality and adapter trimming were performed on the

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no:
---	--------------------------	-----------------------------------	-------	-------	-----------	---------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

raw sequencing reads using Trim Galore! 0.6.3 (35). The reads were mapped to both the human genome (UCSC hg19 with GENCODE annotation) and the SARS-CoV-2 genome (NCBI Assembly ASM985889v3 with Ensembl annotation) using RNA STAR 2.7.5b (36). The resulting mapped reads from each sample were counted by featureCounts 1.6.4 (37) to generate per gene read counts. The raw counts were analyzed for differential expression between experimental conditions using DESeq2 1.22.1 (38), which also generated a normalized gene expression matrix and a PCA plot of the samples.

The number of alternatively spliced XBP1 reads were counted by Integrative Genomics Viewer 2.9.4 (39) using aligned reads data from RNA STAR (see above). The total number of XBP1 reads were counted by featureCounts as above. For each sample, the relative XBP1 splicing was determined by dividing the reads containing the alternative splicing site by the total XBP1 reads.

qRT-PCR

cDNA was synthesized from RNA samples using the High-Capacity cDNA Reverse Transcription Kit (ThermoFisher, 4368813). cDNA samples were diluted in molecular biology grade water, and qRT-PCR experiments were performed on a Roche LightCycler 96 Instrument using the Applied Biosystems PowerUp SYBR Green Master Mix (ThermoFisher, A25776). Results were analyzed by the Roche LightCycler 96 Software. RPL13A was used as a reference gene. The following primer pairs were used (gene name: forward primer, reverse primer):

ERN1: CCGAACGTGATCCGCTACTTCT, CGCAAAGTCCTTCT-GCTCCACA.

EIF2AK3: GTCCCAAGGCTTTGGAATCTGTC, CCTACCAAGA-CAGGAGTTCTGG.

ATF6: CAGACAGTACCAACGCTTATGCC, GCAGAATC-CAGGTGCTTGAAG.

IFIT1: GCCTTGCTGAAAGTGTGGAGGAA, ATCCAGGCGA-TAGGCAGAGATC.

IFIT3: CCTGGAATGCTTACGGCAAGCT, GAGCATCTGA-GAGTCTGCCAA.

ISG15: CTCTGAGCATCCTGGTGAGGAA, AAGGTCAGC-CAGAACAGGTCGT.

OAS1: AGGAAAGGTGCTTCCGAGGTAG, GGAAGTGAAGACAACCAGGT.

SOCS1: TTCGCCCTTAGCGTGAAGATGG, TAGTGCTCCAG-CAGCTCGAAGA.

Alt. spliced XBP1: GCTGAGTCCGCAGCAGGT, CTGGGTC-CAAGTTGTCCAGAAT.

Total XBP1: TGAAAACAGAGTAGCAGCTCAGA, CCAAG-CGCTGTCTTAATC.

RPL13A: CTCAAGGTGTTTGACGGCATCC, TACTTCCAG-CCAACCTCGTGAG.

Clustering of variable genes

The top 5,000 most variable genes were selected, and the normalized gene expression data were analyzed by the Morpheus software (<https://software.broadinstitute.org/morpheus>). K-means clustering with 6 clusters was applied to the gene expression data of the RNA-seq experiment involving CBD and SARS-CoV-2, and K-means clustering with 5 clusters was applied to the gene expression data of the RNA-seq experiment involving CBDV and SARS-CoV-2. For each gene, the normalized expression values of all samples were transformed by subtracting the mean and dividing by the standard

deviation. The transformed gene expression values were used to generate the heatmap.

XBP1 splicing assay

qRT-PCR was used to quantify relative expression of spliced version of XBP1 (XBP1s) by using specific pairs of primers for human alternatively spliced XBP1 and total XBP1 (primer sequences are described above) as previously described (40). Relative percentage of alternative splicing of XBP1 (%XBP1s) was indicated by calculating the ratio of signals between XBP1s and total XBP1.

Ingenuity Pathway Analysis

Expression data (\log_2 fold-change) and predicted activation status of genes were overlaid onto the interferon signaling pathway and the endoplasmic reticulum stress pathway maps using Ingenuity Pathway Analysis. Figures were generated through the use of IPA (QIAGEN Inc). Normalized gene expression values or fold-change (\log_2) of genes were analyzed by the Morpheus software. For each gene, the normalized expression values of all samples were transformed by subtracting the mean and dividing by the standard deviation. The transformed gene expression values were used to generate the heatmaps. IPA-predicted activation z-scores of relevant pathways from the RNA-seq data were also graphed by the Morpheus software.

Gene set enrichment analyses

To identify themes across the 6 clusters, functional gene set enrichment analyses for the genes in each cluster were performed using Metascape (41). The following categories were selected for the enrichment analyses: GO Molecular Functions, KEGG Functional Sets, GO Biological Processes, Canonical Pathways, and KEGG Pathway. Additional parameters for Metascape: Min Overlap = 3, p-value Cutoff = 0.05, Min Enrichment = 1.5. To identify gene sets which activities were reversed by CBD with viral infection, the input gene list includes genes significantly down-regulated by the virus (differential expression comparing veh-infect vs veh-mock, q-value cutoff 0.01) while also significantly up-regulated by CBD (differential expression comparing CBD-infect vs veh-infect, q-value cutoff 0.01). A second list includes genes significantly up-regulated by the virus (differential expression comparing veh-infect vs veh-mock) while also significantly down-regulated by CBD (differential expression comparing CBD-infect vs veh-infect). Gene set enrichment analyses were performed on these two lists of genes using the same Metascape method. The same analyses were also performed on the differential expression data from RNA-seq experiments involving CBDV and SARS-CoV-2 with a q-value cutoff of 0.05. GSEA v4.1.0 was used to perform specific gene set enrichment analyses on Gene Ontology terms PERK-Mediated Unfolded Protein Response and IRE1-Mediated Unfolded Protein Response using the differential expression data from the RNA-seq experiment involving CBD and SARS-CoV-2 (42, 43).

CBD treatment and SARS-CoV-2 challenge in mice

Nine-to-eleven-week-old female K18-hACE2 mice (27) were purchased from Jackson Laboratory (stock #034860). Following acclimation, mice received CBD treatment (20 or 80 mg/kg) via twice daily intraperitoneal injection in a volume of 0.1 ml. The injection solution was prepared immediately before each treatment. First, the CBD powder from Supplier D was dissolved in 100% ethanol. Then the CBD solution was mixed with Cremophor EL (Millipore Sigma 238470) followed by PBS solution at a ratio of 1:1:18. The vehicle

MS no:	Emp name:	Date / Time:	PE's:	AA's:	Comments:	Art no:
RAabi6110/IMMUNOLOGY, LIFE SCIENCES	jrnindesign	1-20-2022 / 22:52				

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

injection solution was prepared by mixing 100% ethanol, Cremophor EL, and PBS at 1:1:18. For each injection, the final amount of CBD was either 20 mg or 80 mg per kg of mouse body weight depending on treatment group. Control groups were treated with vehicle only or received no treatment. Following seven days of treatment, all animals were anesthetized and challenged with 2×10^4 pfu of SARS-CoV-2 (nCoV/Washington/1/2020) via intranasal instillation in a volume of 0.05 ml. After challenge, CBD treatment continued twice daily for an additional 4 days. Mice were also monitored twice daily for the development of clinical disease. Body weights were measured once daily. Five days following virus challenge, all animals were humanely euthanized, and the nasal turbinate and lung tissue collected. Tissues were homogenized in sterile PBS using a hand-held tissue homogenizer (Omni International) and stored at -80°C for virus titration.

SARS-CoV-2 virus titration from mouse tissues by TCID₅₀ assay

Vero E6 cells (ATCC # CRL-1586) were seeded at a density of 20,000 cells/well into 96-well flat-bottomed tissue culture plates (Nunc) and incubated overnight at 37°C with 5% CO_2 and humidity. Homogenized tissues were centrifuged at 8,000 rpm for 10 min at 4°C and the supernatant collected and serially diluted ten-fold (up to 10^{-7}) in viral growth medium (DMEM containing 5% FBS, and 1% antibiotic/antimycotic solution). After overnight incubation, the cell plates were washed twice with PBS and the serial dilutions added to each well in quadruplicate. The plates were further incubated at 37°C in a humidified incubator with 5% CO_2 . After 3 days, cells were stained with 0.1% crystal violet containing 10% neutral-buffered formalin and scored for cytopathic effect (CPE) development. The TCID₅₀ dose was calculated as per Reed and Muench method (44) and corrected for per gram weight of each lung homogenate. All animal work was approved by the University of Louisville Institutional Animal Care and Use Committee. All work with live SARS-CoV-2 was approved by the University Institutional Biosafety Committee and conducted within biosafety level three containment.

Analysis of patient data

All patient data analysis was approved by the National COVID Cohort Collaborative and the University of Chicago Biological Sciences Division institutional review board (IRB21-0591), which granted a waiver of consent because the identities of the study participants cannot readily be ascertained by the investigators, the investigators do not contact the participants, and the investigators will not reidentify participants. A detailed description of the patient data analysis methods and findings is in the Patient Analysis Supplement.

Statistical analysis

Data are shown as means \pm SD. For RNA-seq differential expression analysis, DESeq2 version 1.22.1 was used with a minimum FDR-corrected *P* value (Q value) significance threshold of 0.01 for the RNA-seq experiment involving CBD and SARS-CoV-2, and a threshold of 0.05 for the RNA-seq experiment involving CBDV and SARS-CoV-2. For gene set enrichment analysis, Metascape was used with a minimum *P* value significance threshold of 0.05. For EC50 calculations of drug treatments, GraphPad Prism software was used with a nonlinear curve fit with four parameters. Prism was also used for unpaired *t* tests and one-way ANOVA with statistical significance defined as $p < 0.05$. For the patient data statistical

analysis methods, please refer to the Statistical Analysis section of the Patient Analysis Supplement.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <https://science.org/doi/10.1126/sciadv.abi6110>

[View/request a protocol for this paper from Bio-protocol.](#)

REFERENCES AND NOTES

- S. R. Weiss, J. L. Leibowitz, Coronavirus pathogenesis. *Adv. Virus Res.* **81**, 85–164 (2011).
- S. E. Galloway, P. Paul, D. R. MacCannell, M. A. Johansson, J. T. Brooks, A. M. Neil, R. B. Slayton, S. Tong, B. J. Silk, G. L. Armstrong, M. Biggerstaff, V. G. Dugan, Emergence of SARS-CoV-2 B.1.1.7 lineage - United States, December 29, 2020–January 12, 2021. *MMWR Morb Mortal Wkly Rep* **70**, 95–99 (2021).
- M. Hoffmann, H. Kleine-Weber, S. Schroeder, N. Krüger, T. Herrler, S. Erichsen, T. S. Schiergens, G. Herrler, N.-H. Wu, A. Nitsche, M. A. Müller, C. Drosten, S. Pöhlmann, SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* **181**, 271–280.e8 (2020).
- J. Jaimes, J. Millet, G. Whittaker, Proteolytic cleavage of the SARS-CoV-2 spike protein and the role of the novel S1/S2 Site. *iScience* **23**, 101212 (2020).
- S. Matsuyama, N. Nao, K. Shirato, M. Kawase, S. Saito, I. Takayama, N. Nagata, T. Sekizuka, H. Katoh, F. Kato, M. Sakata, M. Tahara, S. Kutsuna, N. Ohmagari, M. Kuroda, T. Suzuki, T. Kageyama, M. Takeda, Enhanced isolation of SARS-CoV-2 by TMPRSS2-expressing cells. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 7001–7003 (2020).
- W. Wang, C. R. Simoneau, J. Kulsuptrakul, M. Bouhaddou, K. A. Travisano, J. M. Hayashi, J. Carlson-Stevermer, J. R. Zengel, C. M. Richards, P. Fozouni, J. Ok, L. Rodriguez, B. Joehnk, K. Walcott, K. Holden, A. Sil, J. E. Currence, N. J. Krogan, M. Ott, Genetic screens identify host factors for SARS-CoV-2 and common cold coronaviruses. *Cell* **184**, 106–119.e14 (2021).
- W. M. Schneider, J. M. Luna, H.-H. Hoffmann, F. J. Sánchez-Rivera, A. A. Leal, A. W. Ashbrook, J. L. Pen, I. Ricardo-Lax, E. Michailidis, A. Peace, A. F. Stenzel, S. W. Lowe, M. R. MacDonald, C. M. Rice, J. T. Poirier, Genome-scale identification of SARS-CoV-2 and pan-coronavirus host factor networks. *Cell* **184**, 120–132.e14 (2021).
- Z. Daniloski, T. X. Jordan, H.-H. Wessels, D. A. Hoagland, S. Kasela, M. Legut, S. Maniatis, E. P. Mimitou, L. Lu, E. Geller, O. Danziger, B. R. Rosenberg, H. Phatnani, P. Smibert, T. Lappalainen, B. R. tenOever, N. E. Sanjana, Identification of required host factors for SARS-CoV-2 infection in human cells. *Cell* **184**, 92–105.e16 (2021).
- D. W. Kneller, G. Phillips, H. M. O'Neill, R. Jedrzejczak, L. Stols, P. Langan, A. Joachimiak, L. Coates, A. Kovalevsky, Structural plasticity of SARS-CoV-2 3CL Mpro active site cavity revealed by room temperature X-ray crystallography. *Nat. Commun.* **11**, 3202 (2020).
- J. Osipiuk, S. A. Azizi, S. Dvorkin, M. Endres, R. Jedrzejczak, K. A. Jones, S. Kang, R. S. Kathayat, Y. Kim, V. G. Lisnyak, S. L. Maki, V. Nicolaescu, C. A. Taylor, C. Tesar, Y. A. Zhang, Z. Zhou, G. Randall, K. Michalska, S. A. Snyder, B. C. Dickinson, A. Joachimiak, Structure of papain-like protease from SARS-CoV-2 and its complexes with non-covalent inhibitors. *Nat. Commun.* **12**, 743 (2021).
- F. Shahbazi, V. Grandi, A. Banerjee, J. F. Trant, Cannabinoids and cannabinoid receptors: The story so far. *iScience* **23**, 101301 (2020).
- K. M. Nelson, J. Bisson, G. Singh, J. G. Graham, S. N. Chen, J. B. Friesen, J. L. Dahlin, M. Niemitz, M. A. Walters, G. F. Pauli, The essential medicinal chemistry of cannabidiol (CBD). *J. Med. Chem.* **63**, 12137–12155 (2020).
- K. Sekar, A. Pack, Epidiolex as adjunct therapy for treatment of refractory epilepsy: A comprehensive review with a focus on adverse effects. *F1000Res* **8**, F1000 (2019).
- H. I. Lowe, N. J. Toyang, W. McLaughlin, Potential of cannabidiol for the treatment of viral hepatitis. *Pharm. Res.* **9**, 116–118 (2017).
- L. Taylor, B. Gidal, G. Blakey, B. Tayo, G. Morrison, A phase I, randomized, double-blind, placebo-controlled, single ascending dose, multiple dose, and food effect trial of the safety, tolerability and pharmacokinetics of highly purified cannabidiol in healthy subjects. *CNS Drugs* **32**, 1053–1067 (2018).
- S. M. Anil, N. Shalev, A. C. Vinayaka, S. Nadarajan, D. Namdar, E. Belasov, I. Shoval, K. A. Mani, G. Mechrez, H. Koltai, Cannabis compounds exhibit anti-inflammatory activity in vitro in COVID-19-related inflammation in lung epithelial cells and pro-inflammatory activity in macrophages. *Sci. Rep.* **11**, 1462 (2021).
- K. H. D. Crawford, R. Eguia, A. S. Dingens, A. N. Loes, K. D. Malone, C. R. Wolf, H. Y. Chu, M. A. Tortorici, D. Veelsler, M. Murphy, D. Pettie, N. P. King, A. B. Balazs, J. D. Bloom, Protocol and reagents for pseudotyping lentiviral particles with SARS-CoV-2 spike protein for neutralization assays. *Viruses* **12**, 513 (2020).
- C. Hetz, E. Chevet, S. A. Oakes, Proteostasis control by the unfolded protein response. *Nat. Cell Biol.* **17**, 829–838 (2015).
- R. Dash, M. C. Ali, I. Jahan, Y. A. Munni, S. Mitra, M. A. Hannan, B. Timalina, D. F. Oktaviani, H. J. Choi, I. S. Moon, Emerging potential of cannabidiol in reversing proteinopathies. *Ageing Res. Rev.* **65**, 101209 (2021).

MS no:	Emp name:	Date / Time:	PE's:	AA's:	Comments:	Art no:
RAabi6110/IMMUNOLOGY, LIFE SCIENCES	jrnindesign	1-20-2022 / 22:52				

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

20. J. Bechill, Z. Chen, J. W. Brewer, S. C. Baker, Mouse hepatitis virus infection activates the Ire1/XBP1 pathway of the unfolded protein response. *Adv. Exp. Med. Biol.* **581**, 139–144 (2006).
21. T. S. Fung, D. X. Liu, The ER stress sensor IRE1 and MAP kinase ERK modulate autophagy induction in cells infected with coronavirus infectious bronchitis virus. *Virology* **533**, 34–44 (2019).
22. C. Hetz, K. Zhang, R. J. Kaufman, Mechanisms, regulation and functions of the unfolded protein response. *Nat. Rev. Mol. Cell Biol.* **21**, 421–438 (2020).
23. O. Haller, G. Kochs, F. Weber, The interferon response circuit: induction and suppression by pathogenic viruses. *Virology* **344**, 119–130 (2006).
24. D. Blanco-Melo, B. E. Nilsson-Payant, W.-C. Liu, S. Uhl, D. Hoagland, R. Möller, T. X. Jordan, K. Oishi, M. Panis, D. Sachs, T. T. Wang, R. E. Schwartz, J. K. Lim, R. A. Albrecht, B. R. tenOever, Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell* **181**, 1036–1045.e9 (2020).
25. H. Di, H. Elbahesh, M. A. Brinton, Characteristics of human OAS1 isoform proteins. *Viruses* **12**, (2020).
26. T. A. Tummino, V. V. Rezelj, B. Fischer, A. Fischer, M. J. O'Meara, B. Monel, T. Vallet, Z. Zhang, A. Alon, H. R. O'Donnell, J. Lyu, H. Schadt, K. M. White, N. J. Krogan, L. Urban, K. M. Shokat, A. C. Kruse, A. García-Sastre, O. Schwartz, F. Moretti, M. Vignuzzi, F. Pognan, B. K. Shoichet, Phospholipidosis is a shared mechanism underlying the in vitro antiviral activity of many repurposed drugs against SARS-CoV-2. *bioRxiv* 10.1101/2021.03.23.436648 (2021).
27. P. B. McCray Jr., L. Pewe, C. Wohlford-Lenane, M. Hickey, L. Manzel, L. Shi, J. Netland, P. R. Jia, C. Halabi, C. D. Sigmund, C. D. Robinson, J. H. Saltz, H. Spratt, C. Suver, J. Wilbanks, Lethal infection of K18-hACE2 mice infected with severe acute respiratory syndrome coronavirus. *J. Virol.* **81**, 813–821 (2007).
28. M. A. Haendel, C. G. Chute, T. D. Bennett, D. A. Eichmann, J. Guinney, W. A. Kibbe, P. R. O. Payne, E. R. Pfaff, P. N. Robinson, J. H. Saltz, H. Spratt, C. Suver, J. Wilbanks, A. B. Wilcox, A. E. Williams, C. Wu, C. Blacketer, R. L. Bradford, J. J. Cimino, M. Clark, E. W. Colmenares, P. A. Francis, D. Gabriel, A. Graves, R. Hemadri, S. S. Hong, G. Hripscak, D. Jiao, J. G. Klann, K. Kostka, A. M. Lee, H. P. Lehmann, L. Lingrey, R. T. Miller, M. Morris, S. N. Murphy, K. Natarajan, M. B. Palchuk, U. Sheikh, H. Solbrig, S. Visweswaran, A. Walden, K. M. Walters, G. M. Weber, X. T. Zhang, R. L. Zhu, B. Amor, A. T. Girvin, A. Manna, N. Qureshi, M. G. Kurilla, S. G. Michael, L. M. Portilla, J. L. Rutter, C. P. Austin, K. R. Gersing; the N3C Consortium, The National COVID Cohort Collaborative (N3C): Rationale, design, infrastructure, and deployment. *J. Am. Med. Assoc.* **28**, 427–443 (2021).
29. Centers for Disease Control and Prevention, COVID-19, People at Increased Risk, People with Certain Medical Conditions (2021).
30. Q. Zhou, V. Chen, C. P. Shannon, X. S. Wei, X. Xiang, X. Wang, Z. H. Wang, S. J. Tebbutt, T. R. Kollmann, E. N. Fish, Interferon- α 2b Treatment for COVID-19. *Front. Immunol.* **11**, 1061 (2020).
31. T. Muthumalage, I. Rahman, Cannabidiol differentially regulates basal and LPS-induced inflammatory responses in macrophages, lung epithelial cells, and fibroblasts. *Toxicol. Appl. Pharmacol.* **382**, 114713 (2019).
32. Food and Drug Administration (FDA), Application number 210365Orig1s000," *Clinical Pharmacology and Biopharmaceutics Reviews* (2017).
33. B. C. Sorkin, A. J. Kuzak, G. Bloss, N. K. Fukagawa, F. A. Hoffman, M. Jafari, B. Barrett, P. N. Brown, F. D. Bushman, S. J. Casper, F. H. Chilton, C. S. Coffey, M. G. Ferruzzi, D. C. Hopp, M. Kiely, D. Lakens, J. B. MacMillan, D. O. Meltzer, M. Pahor, J. Paul, K. Pritchett-Corning, S. K. Quinney, B. Rehermann, K. D. R. Setchell, N. S. Sipes, J. M. Stephens, D. L. Taylor, H. Tiriac, M. A. Walters, D. Xi, G. Zappala, G. F. Pauli, Improving natural product research translation: From source to clinical trial. *FASEB J.* **34**, 41–65 (2020).
34. E. Afgan, D. Baker, B. Batut, M. van den Beek, D. Bouvier, M. Čech, J. Chilton, D. Clements, N. Coraor, B. A. Grüning, A. Guerler, J. Hillman-Jackson, S. Hiltmann, V. Jalili, H. Rasche, N. Soranzo, J. Goecks, J. Taylor, A. Nekrutenko, D. Blankenberg, The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* **46**, W537–W544 (2018).
35. F. Krueger, Trim Galore. A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files. 517 (2015).
36. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T. R. Gingeras, STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).
37. Y. Liao, G. K. Smyth, W. Shi, featureCounts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).
38. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
39. J. T. Robinson, H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander, G. Getz, J. P. Mesirov, Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).
40. S. B. Yoon, Y. H. Park, S. A. Choi, H. J. Yang, P. S. Jeong, J. J. Cha, S. Lee, S. H. Lee, J. H. Lee, B. W. Sim, B. S. Koo, S. J. Park, Y. Lee, Y. H. Kim, J. J. Hong, J. S. Kim, Y. B. Jin, J. W. Huh, S. R. Lee, B. S. Song, S. U. Kim, Real-time PCR quantification of spliced X-box binding protein 1 (XBP1) using a universal primer method. *PLoS ONE* **14**, e0219978 (2019).
41. Y. Zhou, B. Zhou, L. Pache, M. Chang, A. H. Khodabakhshi, O. Tanaseichuk, C. Benner, S. K. Chanda, Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat. Commun.* **10**, 1523 (2019).
42. A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, J. P. Mesirov, Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 15545–15550 (2005).
43. C. Hetz, F. Martinon, D. Rodriguez, L. H. Glimcher, The unfolded protein response: Integrating stress signals through the stress sensor IRE1 α . *Physiol. Rev.* **91**, 1219–1243 (2011).
44. L. J. Reed, H. Muench, A simple method of estimating fifty per cent endpoints. *Am. J. Epidemiol.* **27**, 493–497 (1938).
45. O. H. D. S. Institute, ATLAS. (2021).
46. O. M. O. P. (OMOP), OMOP Clinical Data Model v5.3.1, Standardized Derived Elements, Drug Era table (2021).
47. Centres for Disease Control and Prevention, COVID-19: When to Quarantine (CDC, 2021).
48. Text Processing Services Library, Regular Expression Operations (Text Processing Services, 2021).
49. J. J. Beltran-Montoya, T. Herreras-Canedo, A. Arzola-Paniagua, F. Vadillo-Ortega, O. F. Dueñas-García, H. Rico-Olvera, A randomized, clinical trial of ketorolac tromethamine vs ketorolac trometamine plus complex B vitamins for cesarean delivery analgesia. *Saudi J Anaesth* **6**, 207–212 (2012).
50. Centers for Disease Control and Prevention, COVID-19-Hospitalization and Death by Age (CDC, 2021).
51. O. M. O. P. (OMOP), OMOP Clinical Data Model v5.3.1, Clinical Data Tables, Person table. (2021).
52. L. Kompaniyets, A. B. Goodman, B. Belay, D. S. Freedman, M. S. Sucusky, S. J. Lange, A. V. Gundlapalli, T. K. Boehmer, H. M. Blanck, Body mass index and risk for COVID-19-related hospitalization, intensive care unit admission, invasive mechanical ventilation, and Death - United States, March-December 2020. *MMWR Morb. Mortal. Wkly Rep.* **70**, 355–361 (2021).
53. C. B. Weir, A. Jan, in *StatPearls*. (Treasure Island (FL), 2021).
54. F. a. D. Administration, (2020).
55. E. C. Leas, E. M. Hendrickson, A. L. Nobles, R. Todd, D. M. Smith, M. Dredze, J. W. Ayers, Self-reported Cannabidiol (CBD) use for conditions with proven therapies. *JAMA Netw. Open* **3**, e2020977 (2020).
56. M. Heptburn, N. Mullaguri, P. George, S. Hantus, V. Punia, A. Bhimraj, C. R. Newey, Acute symptomatic seizures in critically ill patients with COVID-19: Is there an association? *Neurocrit. Care* **34**, 139–143 (2021).
57. S. Corp., Tabulate Two-way reference manual (2021).
58. D. Ho, K. Imai, G. King, E. Suart, A. Whitworth, N. Greifer, MatchIt: Nonparametric Preprocessing for Parametric Causal Inference (2021).
59. P. C. Austin, Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples. *Stat. Med.* **28**, 3083–3107 (2009).
60. D. Pregibon, Data Analytic Methods for Generalized Linear Models, Dissertation, (1979).
61. L. Blizzard, D. W. Hosmer, Parameter estimation and goodness-of-fit in log binomial regression. *Biom. J.* **48**, 5–22 (2006).
62. J. H. C. R. Center, (2021).
63. National COVID Cohort Collaborative, COVID-19 Pehnotype 3.0 Additional Information (N3C, 2021).

Acknowledgments: Overall. The following reagent was deposited by the Centers for Disease Control and Prevention and obtained through BEI Resources, NIAID, NIH: SARS-Related Coronavirus 2, Isolate USA-WA1/2020, NR-52281. Figure 8A was created using BioRender.com. We thank the members of the SARS-CoV-2 host response team in Chicago for stimulating discussions and support with particular thanks to Julian Solway, Rick Morimoto, Nissim Hay, Anne Sperling, HuanHuan Chen, Raphael Lee, Raymond Roos, Shannon Elf, Alexander Muir, Gokhan Mutlu, Jay Pinto, Steven White, Nickolai Dulin, Ray Moellering, Viswanathan Natarajan, Leonitis Plataniias, Karen Ridge and HuanHuan Chen. We thank Dominique Missiakas for facilitating access to the University of Chicago Howard Taylor Ricketts Facility by providing protocols and trained scientists. We also thank Nicole Rosner and Kathleen Cagney for proposing and facilitating analysis of clinical data, Mark Ratain for consideration of pharmacokinetic issues, and Kazuhiko Igarashi for helpful discussions. We thank the University of Chicago Genomics Facility (RRID:SCR_019196) especially Sandhiya Arun and Pieter Faber, for their assistance with RNA sequencing. Finally, we would like to acknowledge the University of Chicago Vice Provost for Research, Karen Kim, and the Dean of the Biological Sciences Division, Kenneth Polonsky, for their steadfast support.

MS no:	Emp name:	Date / Time:	PE's:	AA's:	Comments:	Art no:
RAabi6110/IMMUNOLOGY, LIFE SCIENCES	jrnindesign	1-20-2022 / 22:52				

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

National COVID Cohort Collaborative (N3C). The patient data analyses described in this publication were conducted with data or tools accessed through the NCATS N3C Data Enclave covid.cd2h.org/enclave and supported by NCATS U24 TR002306. This research was possible because of the patients whose information is included within the data from participating organizations (covid.cd2h.org/dtas) and the organizations (<https://ncats.nih.gov/n3c/resources/data-contribution/data-transfer-agreement-signatories>) and scientists who have contributed to the on-going development of this community resource (28). The project described was supported by the National Institute of General Medical Sciences, 5U54GM104942-04. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. The analysis used only de-identified data (i.e., N3C Data Access Tier 2, described at https://covid.cd2h.org/N3C_governance). We gratefully acknowledge contributions from the following N3C core teams:

- *Principal Investigators:* Melissa A. Haendel, Christopher G. Chute, Kenneth R. Gersing, Anita Walden
- *Workstream, subgroup and administrative leaders:* Melissa A. Haendel, Tellen D. Bennett, Christopher G. Chute, David A. Eichmann, Justin Guinney, Warren A. Kibbe, Hongfang Liu, Philip R.O. Payne, Emily R. Pfaff, Peter N. Robinson, Joel H. Saltz, Heidi Spratt, Justin Starren, Christine Suver, Adam B. Wilcox, Andrew E. Williams, Chunlei Wu
- *Key liaisons at data partner sites*
- *Regulatory staff at data partner sites*
- *Individuals at the sites who are responsible for creating the datasets and submitting data to N3C*
- *Data Ingest and Harmonization Team:* Christopher G. Chute, Emily R. Pfaff, Davera Gabriel, Stephanie S. Hong, Kristin Kostka, Harold P. Lehmann, Richard A. Moffitt, Michele Morris, Matvey B. Palchuk, Xiaohan Tanner Zhang, Richard L. Zhu
- *Phenotype Team (Individuals who create the scripts that the sites use to submit their data, based on the COVID and Long COVID definitions):* Emily R. Pfaff, Benjamin Amor, Mark M. Bissell, Marshall Clark, Andrew T. Girvin, Stephanie S. Hong, Kristin Kostka, Adam M. Lee, Robert T. Miller, Michele Morris, Matvey B. Palchuk, Kellie M. Walters
- *Project Management and Operations Team:* Anita Walden, Yooree Chae, Connor Cook, Alexandra Dest, Racquel R. Dietz, Thomas Dillon, Patricia A. Francis, Rafael Fuentes, Alexis Graves, Julie A. McMurry, Andrew J. Neumann, Shawn T. O'Neil, Usman Sheikh, Andréa M. Volz, Elizabeth Zampino
- *Partners from NIH and other federal agencies:* Christopher P. Austin, Kenneth R. Gersing, Samuel Bozzette, Mariam Deacy, Nicole Garbarini, Michael G. Kurilla, Sam G. Michael, Joni L. Rutter, Meredith Temple-O'Connor
- *Analytics Team (Individuals who build the Enclave infrastructure, help create codesets, variables, and help Domain Teams and project teams with their datasets):* Benjamin Amor, Mark M. Bissell, Katie Rebecca Bradwell, Andrew T. Girvin, Amin Manna, Nabeel Qureshi
- *Publication Committee Management Team:* Mary Morrison Saltz, Christine Suver, Christopher G. Chute, Melissa A. Haendel, Julie A. McMurry, Andréa M. Volz, Anita Walden
- *Publication Committee Review Team:* Carolyn Bramante, Jeremy Richard Harper, Wendy Hernandez, Farrukh M Koraisly, Federico Mariona, Saidulu Mattapally, Amit Saha, Satyanarayana Vedula
- *Data partners with released data (50).* Stony Brook University — U24TR002306 • University of Oklahoma Health Sciences Center — U54GM104938: Oklahoma Clinical and Translational Science Institute (OCTSI) • West Virginia University — U54GM104942: West Virginia Clinical and Translational Science Institute (WVCTSI) • University of Mississippi Medical Center — U54GM115428: Mississippi Center for Clinical and Translational Research (CCTR) • University of Nebraska Medical Center — U54GM115458: Great Plains iDeA-Clinical & Translational Research • Maine Medical Center — U54GM115516: Northern New England Clinical & Translational Research (NNE-CTR) Network • Wake Forest University Health Sciences — U1TR001420: Wake Forest Clinical and Translational Science Institute • Northwestern University at Chicago — U1TR001422: Northwestern University Clinical and Translational Science Institute (NUCATS) • University of Cincinnati — U1TR001425: Center for Clinical and Translational Science and Training • The University of Texas Medical Branch at Galveston — U1TR001439: The Institute for Translational Sciences • Medical University of South Carolina — U1TR001450: South Carolina Clinical & Translational Research Institute (SCTR) • University of Massachusetts Medical School Worcester — U1TR001453: The UMass Center for Clinical and Translational Science (UMCCTS) • University of Southern California — U1TR001855: The Southern California Clinical and Translational Science Institute (SC CTSI) • Columbia University Irving Medical Center — U1TR001873: Irving Institute for Clinical and Translational Research • George Washington Children's Research Institute — U1TR001876: Clinical and Translational Science Institute at Children's National (CTSA-CN) • University of Kentucky — U1TR001998: UK Center for Clinical and Translational Science • University of Rochester — U1TR002001: UR Clinical & Translational Science Institute • University of Illinois at Chicago — U1TR002003: UIC Center for Clinical and Translational Science • Penn State Health Milton S. Hershey Medical Center — U1TR002014: Penn State Clinical and Translational Science Institute • The University of Michigan at Ann Arbor — U1TR002240: Michigan Institute for Clinical and Health Research • Vanderbilt University Medical Center — U1TR002243: Vanderbilt Institute for Clinical and Translational Research • University of Washington — U1TR002319: Institute of Translational Health Sciences • Washington University in St. Louis — U1TR002345: Institute of Clinical and Translational Sciences • Oregon Health & Science University — U1TR002369: Oregon Clinical

- and Translational Research Institute • University of Wisconsin-Madison — U1TR002373: UW Institute for Clinical and Translational Research • Rush University Medical Center — U1TR002389: The Institute for Translational Medicine (ITM) • The University of Chicago — U1TR002389: The Institute for Translational Medicine (ITM) • University of North Carolina at Chapel Hill — U1TR002489: North Carolina Translational and Clinical Science Institute • University of Minnesota — U1TR002494: Clinical and Translational Science Institute • Children's Hospital Colorado — U1TR002535: Colorado Clinical and Translational Sciences Institute • The University of Iowa — U1TR002537: Institute for Clinical and Translational Science • The University of Utah — U1TR002538: UHealth Center for Clinical and Translational Science • Tufts Medical Center — U1TR002544: Tufts Clinical and Translational Science Institute • Duke University — U1TR002553: Duke Clinical and Translational Science Institute • Virginia Commonwealth University — U1TR002649: C. Kenneth and Dianne Wright Center for Clinical and Translational Research • The Ohio State University — U1TR002733: Center for Clinical and Translational Science • The University of Miami Leonard M. Miller School of Medicine — U1TR002736: University of Miami Clinical and Translational Science Institute • University of Virginia — U1TR003015: iTHRIVL Integrated Translational health Research Institute of Virginia • Carilion Clinic — U1TR003015: iTHRIVL Integrated Translational health Research Institute of Virginia • University of Alabama at Birmingham — U1TR003096: Center for Clinical and Translational Science • Johns Hopkins University — U1TR003098: Johns Hopkins Institute for Clinical and Translational Research • University of Arkansas for Medical Sciences — U1TR003107: UAMS Translational Research Institute • Nemours — U54GM104941: Delaware CTR ACCEL Program • University Medical Center New Orleans — U54GM104940: Louisiana Clinical and Translational Science (LA CaTS) Center • University of Colorado Denver, Anschutz Medical Campus — U1TR002535: Colorado Clinical and Translational Sciences Institute • Mayo Clinic Rochester — U1TR002377: Mayo Clinic Center for Clinical and Translational Science (CaTS) • Tulane University — U1TR003096: Center for Clinical and Translational Science • Loyola University Medical Center — U1TR002389: The Institute for Translational Medicine (ITM) • Advocate Health Care Network — U1TR002389: The Institute for Translational Medicine (ITM) • OCHIN — INV-018455: Bill and Melinda Gates Foundation grant to Sage Bionetworks
- *Additional data partners who have signed DTA and data release pending (35).* The Rockefeller University — U1TR001866: Center for Clinical and Translational Science • The Scripps Research Institute — U1TR002550: Scripps Research Translational Institute • University of Texas Health Science Center at San Antonio — U1TR002645: Institute for Integration of Medicine and Science • The University of Texas Health Science Center at Houston — U1TR003167: Center for Clinical and Translational Sciences (CCTS) • NorthShore University HealthSystem — U1TR002389: The Institute for Translational Medicine (ITM) • Yale New Haven Hospital — U1TR001863: Yale Center for Clinical Investigation • Emory University — U1TR002378: Georgia Clinical and Translational Science Alliance • Weill Medical College of Cornell University — U1TR002384: Weill Cornell Medicine Clinical and Translational Science Center • Montefiore Medical Center — U1TR002556: Institute for Clinical and Translational Research at Einstein and Montefiore • Medical College of Wisconsin — U1TR001436: Clinical and Translational Science Institute of Southeast Wisconsin • University of New Mexico Health Sciences Center — U1TR001449: University of New Mexico Clinical and Translational Science Center • George Washington University — U1TR001876: Clinical and Translational Science Institute at Children's National (CTSA-CN) • Stanford University — U1TR003142: Spectrum: The Stanford Center for Clinical and Translational Research and Education • Regenstrief Institute — U1TR002529: Indiana Clinical and Translational Science Institute • Cincinnati Children's Hospital Medical Center — U1TR001425: Center for Clinical and Translational Science and Training • Boston University Medical Campus — U1TR001430: Boston University Clinical and Translational Science Institute • The State University of New York at Buffalo — U1TR001412: Clinical and Translational Science Institute • Aurora Health Care — U1TR002373: Wisconsin Network For Health Research • Brown University — U54GM115677: Advance Clinical Translational Research (Advance-CTR) • Rutgers, The State University of New Jersey — U1TR003017: New Jersey Alliance for Clinical and Translational Science • Loyola University Chicago — U1TR002389: The Institute for Translational Medicine (ITM) • #N/A — U1TR001445: Langone Health's Clinical and Translational Science Institute • Children's Hospital of Philadelphia — U1TR001878: Institute for Translational Medicine and Therapeutics • University of Kansas Medical Center — U1TR002366: Frontiers: University of Kansas Clinical and Translational Science Institute • Massachusetts General Brigham — U1TR002541: Harvard Catalyst • Icahn School of Medicine at Mount Sinai — U1TR001433: ConduITS Institute for Translational Sciences • Ochsner Medical Center — U54GM104940: Louisiana Clinical and Translational Science (LA CaTS) Center • HonorHealth — None (Voluntary) • University of California, Irvine — U1TR001414: The UC Irvine Institute for Clinical and Translational Science (ICTS) • University of California, San Diego — U1TR001442: Altman Clinical and Translational Research Institute • University of California, Davis — U1TR001860: UC Davis Health Clinical and Translational Science Center • University of California, San Francisco — U1TR001872: UCSF Clinical and Translational Science Institute • University of California, Los Angeles — U1TR001881: UCLA Clinical Translational Science Institute • University of Vermont — U54GM115516: Northern New England Clinical & Translational Research (NNE-CTR) Network • Arkansas Children's Hospital — U1TR003107: UAMS Translational Research Institute

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no:
---	--------------------------	-----------------------------------	-------	-------	-----------	---------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

•*Additional N3C acknowledgments.* We would also like to thank others not listed above: Joy Alamgir (ARIScience), Seth Russell, MS (University of Colorado), and Hui Zhang, PhD MBA (University of Chicago), who created N3C concept sets that we used to define some of the medical condition indicator covariates for our statistical analysis. We would also like to thank Steve Johnson, PhD (University of Minnesota) for writing a BMI calculation Knowledge Object that we modified for calculating BMI values from weight and height data.

Funding: This work was supported directly by: BIG Vision grant from the University of Chicago (M.R.R.). National Institutes of Health grant R01 GM121735 (M.R.R.). National Institutes of Health grant R01 CA184494 (M.R.R.). National Institutes of Health grant R01 AI137514 (G.R.). National Institutes of Health grant R01 AI127518 (G.R.). National Institutes of Health grant R01 AI134980 (G.R.). National Institutes of Health grant R01 CA219815 (S.A.O.). National Institutes of Health grant R35 GM119840 (B.C.D.). National Institutes of Health grant P30 CA014599 (University of Chicago Comprehensive Cancer Center Support grant). Harry B. and Leona M. Helmsley Charitable Trust (K. E. P). The patient analysis component of this work was supported indirectly by the funding listed in the N3C section of the Acknowledgements section. **Author contributions:** Conceptualization: MRR, LCN, DY, GR, SAO, GFP, DOM, ND, TJB. Methodology: MRR, LCN, DY, GR, SAO, DOM, GFP, ND, BCD, TJB. Software: DY, TJB. Formal analysis: DY,

TJB. Investigation: LCN, DY, VN, TJB, TO, SN-C, JBF, ND, AM, CD, Diane Silva, HG, LS, LR-M, AV, ES, KAJ, S-AA. Resources: GR, MRR, GFP, ST, BCD. Data Curation: DOM, TJB, DY, National COVID Cohort Collaborative Consortium. Writing – original draft: MRR. Writing – review & editing: LCN, DY, VN, TJB, SN-C, JBF, ND, AM, KAJ, Diane Silva, JMM, BCD, ST, SAO, GFP, DOM, GR, MRR. Visualization: LCN, DY, ND, AM, SN-C, BCD, JBF. Supervision: MRR, GR, DOM, GFP, SAO, ST, BCD. Project Administration: MRR. Funding acquisition: MRR. For the animal experiment: Investigation: Divyasha Saxena, CDA. Formal analysis: Divyasha Saxena, JDG. Methodology: JDG, JKD, Divyasha Saxena, WES, KEP. Obtaining approvals: WES, JDG. Funding acquisition: KEP. **Competing interests:** S.A.O. is a cofounder and consultant at OptiKira., L.L.C. (Cleveland, OH). The authors declare no other competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper, the Supplementary Materials, and/or the Supplementary Data S1. Raw and processed RNA-seq data were deposited into the GEO database (GSE168797).

Submitted 19 March 2021
 Accepted 29 December 2021
 Published First Release 20 January 2022
 10.1126/sciadv.abi6110

MS no: RAabi6110/IMMUNOLOGY, LIFE SCIENCES	Emp name: jrnindesign	Date / Time: 1-20-2022 / 22:52	PE's:	AA's:	Comments:	Art no:
---	--------------------------	-----------------------------------	-------	-------	-----------	---------

Teaser: Cannabidiol from the cannabis plant has potential to prevent and inhibit SARS-CoV-2 infection.

Cannabidiol inhibits SARS-CoV-2 replication through induction of the host ER stress and innate immune responses

Long Chi NguyenDongbo YangVlad NicolaescuThomas J. BestHaley GulaDivyasha SaxenaJon D. GabbardShao-Nong ChenTakashi OhtsukiJohn Brent FriesenNir DraymanAdil MohamedChristopher DannDiane SilvaLydia Robinson-MailmanAndrea ValdespinoLeticia StockEva SuárezKrysten A. JonesSaara-Anne AziziJennifer K. DemarcoWilliam E. SeversonCharles D. AndersonJames Michael MillisBryan C. DickinsonSava# TayScott A. OakesGuido F. PauliKenneth E. PalmerThe National COVID Cohort Collaborative ConsortiumDavid O. MeltzerGlenn RandallMarsha Rich Rosner

Sci. Adv., Ahead of Print

View the article online

<https://www.science.org/doi/10.1126/sciadv.abi6110>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of think article is subject to the [Terms of service](#)